arXiv:1204.0488v1 [math.AG] 2 Apr 2012

NONCOMMUTATIVE DESINGULARIZATION OF ORBIT CLOSURES FOR SOME REPRESENTATIONS OF GL_n

JERZY WEYMAN AND GUFANG ZHAO

Abstract. We describe noncommutative desingularizations of determinantal varieties, determinantal varieties defined by minors of generic symmetric matrices, and pfaffian varieties defined by pfaffians of generic anti-symmetric matrices. For maximal minors of square matrices and symmetric matrices, this gives a non-commutative crepant resolution. Along the way, we describe a method to calculate the quiver with relations for any non-commutative desingularizations coming from exceptional collections over partial flag varieties.

Contents

1. Introduction 1
2. Constructions and formulas for non-commutative desingularizations 6
3. Grassmannians 14
4. Equivariant quivers 19
5. Determinantal varieties of symmetric matrices 22
6. Pfaffian varieties of anti-symmetric matrices 31
7. Determinantal varieties 33
8. Other examples 36
References 39

1. Introduction

For a singular variety X, a *non-commutative desingularization* (Definition 2.2) is a coherent sheaf of associative algebras $\mathscr{A}$ over a proper scheme Y over X birationally equivalent to X, such that $\mathscr{A}$ is generically a sheaf of matrix algebras and has finite homological dimension. This notion arises from the study of the derived categories of coherent sheaves, as well as theoretical physics. In the case that $X = \operatorname{Spec} R$ for a Gorenstein ring R, Van den Bergh introduced a notion of *non-commutative crepant resolution* (Definition 2.4, c.f. also [V04b]), which means a non-commutative desingularization whose global section is maximal Cohen-Macaulay. This notion is a generalization of an existent notion of crepant resolution, and is introduced with the hope that any two crepant resolutions, commutative or not, are derived equivalent.

The idea of non-commutative desingularization can be traced back to the general theory of derived Morita equivalence due to Rickard and others. Let k be a field. An ordered set of

---

Date: March 2, 2013.
2000 *Mathematics Subject Classification.* Primary 14F05, 4E05, 16G20; Secondary 14A22, 13D02, 14L35 .
Key words and phrases. determinantal variety, symmetric minor, pfaffian, partial flag variety, exceptional collection, derived equivalence, quiver with relations.

objects $\nabla = \{\nabla_\alpha, \alpha \in I\}$ in a triangulated k-linear category D is called (strong) exceptional if we have $\mathrm{Ext}^\bullet(\nabla_\alpha, \nabla_\beta) = 0$ for $\alpha < \beta$; $\mathrm{Ext}^n(\nabla_\alpha, \nabla_\alpha) = 0$ for $n \neq 0$, and $\mathrm{End}(\nabla_\alpha) = k$. An exceptional set is said to be full if it generates D. If X is a projective variety, $(E_0, \cdots, E_n)$ is a full exceptional collection in the derived category of coherent sheaves on X. Then there is an equivalence of derived categories

$$R\,\mathrm{Hom}(\oplus_i E_i, -) : D^b(X) \to D^b(A\text{-}mod),$$

where A-*mod* is the category of finitely generated left modules over $A := \mathrm{End}(\oplus_i E_i)$ with the opposite multiplication. Note that by properness of X, the algebra A is finite dimensional and can be described as the path algebra of a quiver with relations.

Let Z be a quasi-projective variety, e.g., the total space of a vector bundle over a projective variety X, Z rarely admits an exceptional collection. Nevertheless, if Z is projective over some affine scheme of finite type, it admits tilting objects. And in this case Z is derived equivalent to the endomorphism algebra of a tilting object, as has been explained in [HV07] and will be reviewed in Section 2. Here by a tilting object, we mean a perfect complex T in $D^b(Z)$ such that $\mathrm{Ext}^i(T, T) = 0$ for all $i \neq 0$, and T generates the entire derived category, in the sense that the smallest triangulated subcategory containing T closed under direct summands is the entire derived category. In all cases we are interested in, the tilting object can always be chosen as a vector bundle, considered as a complex of coherent sheaves concentrated on degree 0. If this happens, the assumption that T generates $D^b(Z)$ can be replaced by that $\mathrm{Ext}^\bullet(T, C) = 0$ for some complex C implies C is exact (see, e.g. [BH10]). But the fact that Z is non-compact makes the endomorphism algebra more complicated than the proper case.

If X is a smooth projective Fano variety and $Z = \omega_X$ is the total space of its canonical bundle, the inverse images of some exceptional collections on X to ω_X has been used by Tom Bridgeland and others to study t-structures in the derived categories of coherent sheaves on ω_X (see, e.g. [Br05]).

Let G be a reductive group and P a parabolic subgroup. Let $p : Z \to G/P$ be an equivariant vector bundle. In this paper and subsequent ones, we use the inverse image of the exceptional collections on G/P to Z, with special emphasis on the situation when the total space Z is a commutative desingularization of a orbit closure of some representations. When the vector bundle is the commutative desingularization of generic determinantal variety defined by maximal minors, a non-commutative desingularization of this nature has been studied by R. Buchweitz, G. Leuschke, and M. van den Bergh in [BLV09]. In the current paper, we describe separately the quiver with relations of noncommutative desingularizations of higher corank determinantal varieties, symmetric determinantal varieties, and anti-symmetric determinantal varieties.

The study of exceptional collections was initiated by Beilinson and Kapranov who dealt with the case of the projective spaces and partial flag varieties. According to the result of Kapranov, the set

$$\{\mathbb{S}_\alpha \mathcal{Q}^* \mid \alpha \subset ((n-r)^r)\}$$

is an exceptional collection over the grassmannian $\mathrm{Grass}_{n-r}(E)$ of $(n-r)$-planes in E, where $\mathbb{S}_\lambda$ is the Schur functor applied to the partition λ, the index set is the set of all subpartitions of $((n-r)^r)$, the square with $n-r$ columns and r rows, and $\mathcal{Q}$ is the tautological rank r quotient bundle over $\mathrm{Grass}_{n-r}(E)$.

Let $p : Z \to G/P$ be an equivariant vector bundle, and let $\nabla(G/P) = \{\nabla_\alpha \mid \alpha \in I\}$ be a full exceptional collection on G/P. On the total space Z, the inverse image $p^*(\oplus_\alpha \nabla_\alpha)$ can fail to be a tilting object. Moreover, its endomorphism algebra $\Lambda := \mathrm{End}_{\mathscr{O}_Z}(p^*(\oplus_\alpha \nabla_\alpha))$ is usually infinite dimensional as a vector space.

It is easy to show (Proposition 2.7) that for any tilting bundle $\mathscr{T}il$ on G/P, (in particular for $\oplus_\alpha \nabla_\alpha$,) the inverse image $p^*\mathscr{T}il$ is a tilting bundle on Z if the $p^*\mathscr{E}nd_{\mathscr{O}_{G/P}}(\mathscr{T}il)$ has no higher cohomology. If $q : Z \to \operatorname{Spec} R$ is a commutative desingularization of an affine (integral) variety $\operatorname{Spec} R$, and if the vector bundle Z has rank higher than 1, then we show that $\mathrm{End}_{\mathscr{O}_Z}(p^*\mathscr{T}il) \cong \mathrm{End}_R(q_*p^*\mathscr{T}il)$ and $\mathrm{End}_R(q_*p^*\mathscr{T}il)$ is a non-commutative desingularization of $\operatorname{Spec} R$.

In the case of homogenenous space G/P the exceptional collection often consists of equivariant vector bundles. In such case the objects in $D^b(Z)$ corresponding to the simples are obtained by pushing forward the dual exceptional collection (Definition 2.12) over G/P.

Let $\mathfrak{A}$ be a k-linear category. Recall that for an exceptional collection ∇, the *dual collection* $\Delta = \{\Delta_\alpha \mid \alpha \in I\}$ is another subset of objects in $D^b(\mathfrak{A})$, in bijection with ∇, such that $\mathrm{Ext}^\bullet(\Delta_\beta, \nabla_\alpha) = 0$ for $\beta > \alpha$, and there exists an isomorphism $\Delta_\beta \cong \nabla_\beta \bmod D_{<\beta}$, where $D_{<\alpha}$ is the full triangulated subcategory generated by $\{\nabla_\beta \mid \beta < \alpha\}$.

Let $\Lambda = \mathrm{End}_{\mathscr{O}_Z}(p^*(\oplus_\alpha \nabla_\alpha))$ and $S_\beta = R\,\mathrm{Hom}_{\mathscr{O}_Z}(p^*(\oplus_\alpha \nabla_\alpha), u_*\Delta_\beta)$, where $u : G/P \to Z$ is the zero section.

We prove

Theorem A (Theorem 2.25)**.** Let $\nabla(G/P) = \{\nabla_\alpha \mid \alpha \in I\}$ be a full exceptional collection consisting of equivariant sheaves over G/P with the dual collection Δ. Assume the resolution $q : Z \to \operatorname{Spec} R$ is G-equivariant with $q^{-1}(0) = G/P$, and the only fixed closed point of $\operatorname{Spec} R$ is $\{0\} \subset \operatorname{Spec} R$. Assume moreover that $p^*(\oplus_\alpha \nabla_\alpha)$ is a tilting bundle over Z, $\Lambda \cong \mathrm{End}_R(q_*p^*(\oplus_\alpha \nabla_\alpha))$, and $\mathrm{End}_R(q_*p^*(\oplus_\alpha \nabla_\alpha))$ is a non-commutative desingularization. Then,

(1) S_α's are equivariant simple objects in Λ-*mod*;
(2) a basis of the vector space $\mathrm{Ext}^1_\Lambda(S_\alpha, S_\beta)^*$ generates Λ over $\oplus_\alpha \mathbb{C}_\alpha$;
(3) with the generators for $\mathrm{End}_R(q_*p^*\mathscr{T}il)$ as above, $\mathrm{Ext}^2_{\mathscr{O}_Z}(\Delta_i, \Delta_j)^*$ generates the relations.

If $\operatorname{Spec} R$ is the closure of an orbit in a representation of G, and the commutative desingularization is an equivariant vector bundle over G/P, the above theorem and the discussions proceeding it give a general approach to calculate a non-commutative desingularization of orbit closures. The drawback of our approach, compared to [BLV09] and [BLV11], is that our approach does not work well in positive characteristic, although a fair part of Section 2 holds in a characteristic free fashion. Unless otherwise specified, **we work over a field of characteristic zero**. We will write the field as $\mathbb{C}$ if its characteristic is zero, and k if its characteristic is arbitrary.

The non-commutative desingularization $\mathrm{End}_R(q_*p^*\mathscr{T}il)$, expressed by idempotents, generators and relations as above, can also be expressed as the path algebra of a quiver with relations. Note that the vector space spanned by the set of arrows between any two vertices is naturally a representation of G. We find it convenient to introduce the language of equivariant quivers, as all the Ext's are naturally representations of G. For the precise definitions of equivariant quivers and their representations, see Section 4. An *equivariant quiver* is a triple $Q = (Q_0, Q_1, \alpha)$, where (Q_0, Q_1) is a quiver and α is an assignment associating each arrow $q \in Q_1$ a finite dimensional irreducible representation of G. For any equivariant quiver, there is an underlying usual quiver, upon choosing a basis for each representation associated to each arrow. The path algebra of an equivariant quiver is endowed with a natural rational G-action, so that we can consider the equivariant representations of it. While imposing relations to an equivariant quiver, we require the relations to be subrepresentations. In fact, if the equivariant quiver gives the endomorphism ring of a tilting object over G/P, the derived category of equivariant sheaves over G/P is equivalent to the derived category of equivariant representations of the path algebra with relations of the equivariant quiver.

Let $\operatorname{Spec} R^s$ be a GL_n orbit closure in the space of $n \times n$ symmetric matrices, we do get a non-commutative desingularization by pulling-back the Kapranov's exceptional collection.[1] More precisely, we assume E is a vector spaces over k of dimension n and H^s be the affine space $\operatorname{Sym}_2(E^*)$. Upon choosing a set of basis for E, the coordinate ring of H^s can be identified with $S^s = k[x_{ij}]_{i \leq j}$. Then R^s is the quotient of S^s by the ideal generated by the $(r+1) \times (r+1)$ minors of the generic matrix (x_{ij}).

Let $0 \to \mathcal{R} \to E \times \text{Grass} \to Q \to 0$ be the tautological sequence over Grass, where Grass is the grassmannian of $(n-r)$-planes in E. Then the total space $\mathcal{Z}^s$ of $\operatorname{Sym}_2 Q^*$ is a commutative desingularization of $\operatorname{Spec} R^s$, as is proved in [W03].

We consider the pull back of $\mathscr{T}il_K = \oplus_{\alpha \subset ((n-r)^r)} \mathbb{S}_\alpha Q^*$, the Kapranov's tilting bundle over $\text{Grass}_{n-r}(E)$ by $p' : \mathcal{Z}^s \to \text{Grass}$.

Theorem B (Proposition 5.3, Proposition 5.6)**.** *Let $\mathscr{T}il_K$ be the Kapranov's tilting bundle over $\text{Grass}_{n-r}(E)$, and $\mathcal{Z}^s$ is the total space of $\operatorname{Sym}_2 Q^*$ which desingularizes the rank r determinantal variety $\operatorname{Spec} R^s$ of symmetric matrices.*

1. *The bundle $p'^* \mathscr{T}il_K$ is a tilting bundle over $\mathcal{Z}^s$, i.e., $\operatorname{Ext}^i_{\mathcal{Z}^s}(p'^* \mathscr{T}il_K, p'^* \mathscr{T}il_K) = 0$ for $i > 0$ and $\operatorname{Ext}^\bullet(p'^* \mathscr{T}il_K, C) = 0$ implies C is an exact complex. In particular, $D^b(Coh(\mathcal{Z}^s)) \cong D^b(\operatorname{End}_{\mathcal{Z}^s}(p'^* \mathscr{T}il_K)\text{-}mod)$.*
2. *The map $\operatorname{End}_{\mathcal{Z}^s}(p'^* \mathscr{T}il_K) \to \operatorname{End}_S(q'_* p'^* \mathscr{T}il_K)$ is an isomorphism of R^s-algebra, and this algebra is a non-commutative desingularization of $\operatorname{Spec} R^s$. In other words, it has finite global dimension. Moreover, if $r = n-1$, it is maximal Cohen-Macaulay over R^s.*

Now we describe the equivariant quiver with relations for the non-commutative desingularization. We will use $C^\beta_{\alpha,\mu}$ for the Littlewood-Richardson coefficient. For any two vertices α and β, the space of paths from β to α will be denoted by $\operatorname{Hom}(\beta, \alpha)$.

The equivariant quiver with relations of the endomorphism ring $\operatorname{End}_S(q'_* p'^* \mathscr{T}il_K)$ is given as follows (see Proposition 5.14 and Proposition 5.15). The set of vertices is indexed by subpartitions of $((n-r)^r)$.

- In the case $n-r = 1$, the set of arrows from β to α is given by E if $C^\alpha_{\beta,(1,0,\cdots,0)} \neq 0$ or $C^\beta_{\alpha,(1,0,\cdots,0)} \neq 0$. No arrows otherwise. The relations are generated by the following representations in the space $\operatorname{Hom}(\beta, \alpha)$
$$(C^\alpha_{\beta,(1,1,0,\cdots,0)} \mathbb{S}_2 E) \oplus (C^\beta_{\alpha,(1,1,0,\cdots,0)} \mathbb{S}_2 E) \oplus (\delta^\beta_\alpha \wedge^2 E).$$
- In the case $n-r = 2$, arrows from β to α is given by E if $C^\alpha_{\beta,(1,0,\cdots,0)} \neq 0$ and $\mathbb{C}$ if $C^\beta_{\alpha,(1,1,0,\cdots,0)} \neq 0$. No arrows otherwise. The relations are generated by the following representations in the group $\operatorname{Hom}(\beta, \alpha)$
$$(C^{\beta^t}_{\alpha^t,(1,-1)} \wedge^2 E) \oplus (C^{\beta^t}_{\alpha^t,(-1,-2)} E) \oplus (C^\beta_{\alpha,(1,1,0,\cdots,0)} \mathbb{S}_2 E) \oplus (C^\beta_{\alpha,(2,0,\cdots,0)} \wedge^2 E).$$
- In the case $n-r \geq 3$, The set of arrows has the same description as in the case $n-r = 2$. The relations are generated by the following representations in the group

---

[1] We use the upper script s to remind us that we are in the symmetric matrix case. Similarly, later on we will use the upper script a in the skew-symmetric matrix case.

$\operatorname{Hom}(\beta, \alpha)$

$$(C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-1,-2)}\mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-1,-1)}E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)}\mathbb{S}_2 E) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E).$$

Now let $\operatorname{Spec} R^a$ be a GL_n-orbit closure in the space of $n \times n$ skew-symmetric matrices, we also get a non-commutative desingularization by pulling-back the Kapranov's exceptional collection. Let $0 \to \mathcal{R} \to E \times \mathrm{Grass} \to \mathcal{Q} \to 0$ be the tautological sequence over Grass, where Grass is the grassmannian of $(n-r)$-planes in E. Then the total space $\mathcal{Z}^a$ of $\wedge^2 \mathcal{Q}^*$ is a commutative desingularization of $\operatorname{Spec} R^a$, as is proved in [W03].

We consider the pull back of the Kapranov's tilting bundle $\mathcal{T}il_K = \oplus_{\alpha \subset ((n-r)^r)} \mathbb{S}_\alpha \mathcal{Q}^*$ over $\mathrm{Grass}_{n-r}(E)$ by $p' : \mathcal{Z}^a \to \mathrm{Grass}$.

Theorem C (Proposition 6.2, Proposition 6.4)**.** *Let $\mathcal{T}il_K$ be the Kapranov's tilting bundle over $\mathrm{Grass}_{n-r}(E)$, and $\mathcal{Z}^a$ is the total space of $\wedge^2 Q^*$ which desingularizes the rank r Pfaffian variety $\operatorname{Spec} R^a$ of skew-symmetric matrices.*

1. *The bundle $p'^* \mathcal{T}il_K$ is a tilting bundle over $\mathcal{Z}^a$, i.e., $\operatorname{Ext}^i_{\mathcal{Z}^a}(p'^* \mathcal{T}il_K, p'^* \mathcal{T}il_K) = 0$ for $i > 0$ and $\operatorname{Ext}^\bullet(p'^* \mathcal{T}il_K, C) = 0$ implies C is an exact complex. In particular, $D^b(Coh(\mathcal{Z}^a)) \cong D^b(\operatorname{End}_{\mathcal{Z}^a}(p'^* \mathcal{T}il_K)\text{-}mod)$.*
2. *If $r > 2$, the map $\operatorname{End}_{\mathcal{Z}^a}(p'^* \mathcal{T}il_K) \to \operatorname{End}_S(q'_* p'^* \mathcal{T}il_K)$ is an isomorphism of R^a-algebra, and this algebra is a non-commutative desingularization of $\operatorname{Spec} R^a$. In other words, it has finite global dimension.*

The quiver with relations of the endomorphism ring is given as follows (see Proposition 6.7 and Proposition 6.8). The set of vertices is indexed by subpartitions of $((n-r)^r)$.

- In the case $n - r = 1$, arrows from β to α is given by E if $C^{\alpha}_{\beta,(1,0,\cdots,0)} \neq 0$ and $\mathbb{C}$ if $C^{\beta}_{\alpha,(1,1,0,\cdots,0)} \neq 0$. No arrows otherwise. The relations are generated by the following representations in the group $\operatorname{Hom}(\beta, \alpha)$
$$(C^{\beta^t}_{\alpha^t,(1,0,\cdots,0)} \wedge^3 E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)}\mathbb{S}_2 E).$$
- In the case $n - r = 2$, there is one more arrow from β to α given by $\mathbb{C}$ for $C^{\beta^t}_{\alpha^t,(1,-1,0,\ldots,0)}$ besides the above ones. No arrows otherwise. The relations are generated by the following representations in the group $\operatorname{Hom}(\beta, \alpha)$
$$(C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-3)}\mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-2)}E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)}\mathbb{S}_2 E) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E).$$
- In the case $n - r \geq 3$, The set of arrows has the same description as in the case $n - r = 1$. The set of relations has the same description as in the case $n - r = 2$.

In the case of generic determinantal varieties, the quiver with relations are already known to the experts. We used the geometric technique in [W03] to study the decomposition of the non-commutative desingularization as representations.

CONVENTIONS. For an abelian category $\mathfrak{A}$, its derived category will be denoted by $D(\mathfrak{A})$, and its bounded derived category $D^b(\mathfrak{A})$. For a scheme X, we denote the abelian category of quasi-coherent sheaves over it by $Qcoh(X)$, and the abelian category of coherent sheaves $Coh(X)$. For short, $D^b(X) = D^b(Qcoh(X))$. For a ring A, commutative or not, A-*Mod* will be used to denote the abelian category of (left) A-modules, and A-*mod* the abelian category of finitely generated (left) A-modules. For a partition λ, we denote by $\mathbb{S}_\lambda$ the corresponding Schur functor. We will identify partitions and Young diagrams and following the conventions in [F97]. For a vector bundle $\mathcal{F}$ over X, we denote its dual bundle $\mathcal{H}om_X(\mathcal{F}, \mathcal{O})$ by $\mathcal{F}^*$. Similarly, for a vector space E, its dual space is denoted by E^*. For any R-module M, when we talk about the algebra structure of $\operatorname{End}_R(M)$, we consider the opposite multiplication. Similar for $\mathscr{E}nd_{\mathscr{O}_X}(\mathscr{F})$ for any quasicoherent sheaf $\mathscr{F}$ on X.

ACKNOWLEDGEMENTS. Many ideas in this paper are inspired by the work of R. Buchweitz, G. Leuschke, and M. van den Bergh in [BLV09]. The second author would like to thank Yi Zhu for helpful discussions related to the proof of Proposition 2.19, and Roman Bezrukavnikov for confirming Remark 2.18. He is also grateful to Yaping Yang for innumerable times of fruitful discussions.

2. Constructions and formulas for non-commutative desingularizations

2.1. **General notions about tilting bundles.** For a scheme X, a vector bundle (or more generally a perfect complex) $\mathscr{T}il$ is called a *tilting bundle* over X if it satisfies the following:

1. $\mathscr{T}il^{\perp} = 0$ in $D^b(X)$, i.e., for any complex M, $\mathrm{Hom}_{D^b(X)}(\mathscr{T}il, M[i]) = 0$ for all i implies $M = 0$, where $[1]$ is the shifting functor;
2. $\mathrm{Ext}^i(\mathscr{T}il, \mathscr{T}il) = 0$ for $i > 0$.

Theorem 2.1 (7.6 in [HV07])**.** *For X a projective scheme over a Noetherian affine scheme of finite type, and $\mathscr{T}il \in D(Qcoh(X))$ a tilting object. We have the following*

(1) $R\,\mathrm{Hom}_{\mathscr{O}_X}(\mathscr{T}il, -)$ *induces an equivalence*

$$D(Qcoh(X)) \cong D(\mathrm{End}_{\mathscr{O}_X}(\mathscr{T}il)\text{-}Mod).$$

(2) *This equivalence restricts to an equivalence*

$$D^b(Coh(X)) \cong D^b(\mathrm{End}_{\mathscr{O}_X}(\mathscr{T}il)\text{-}mod).$$

(3) *If X is smooth then $\mathrm{End}_{\mathscr{O}_X}(\mathscr{T}il)$ has finite global dimension.*

Definition 2.2. [§ 5 in [BO02]] For an algebraic variety X, a *noncommutative (birational) desingularization* of X is a pair $(p, \mathscr{A})$ consisting of a proper birational morphism $p : Y \to X$ and an algebra $\mathscr{A} = \mathscr{E}nd_{\mathscr{O}_Y}(\mathscr{F})$ on Y, the sheaf of local endomorphisms of a reflexive coherent $\mathscr{O}_Y$-module $\mathscr{F}$, such that the abelian category of sheaves of right modules $\mathscr{A}$ has finite homological dimension.

Remark 2.3. In the original definition of [BO02], the sheaf $\mathscr{F}$ is only required to be torsion free.

Definition 2.4. For a Gorenstein ring R, we call an endomorphism ring $A = \mathrm{End}_R(M)$ for a reflexive R-module M a non-commutative crepant resolution if A has finite global dimension and is maximal Cohen-Macaulay.

2.2. **Preimage of tilting bundles and noncommutative desingularization.** Let G be a reductive group, $P < G$ a parabolic subgroup. Suppose we have a tilting bundle $\mathscr{T}il_{G/P}$ over the partial flag variety G/P, and suppose the total space Z of a vector subbundle of $\mathbb{A}^n \times G/P$ over G/P desingularizes an affine subvariety $\mathrm{Spec}\, R \subseteq \mathbb{A}^n$, i.e., the projection $\mathbb{A}^n \times G/P \to \mathbb{A}^n$ restricted to Z is birational onto $\mathrm{Spec}\, R$, so that the restriction is a desingularization. We seek the conditions for the inverse image of $\mathscr{T}il_{G/P}$ to be a tilting bundle over Z, and to give a non-commutative crepant resolution of $\mathrm{Spec}\, R$.

(1)
$$\begin{tikzcd}
Z \arrow[dr, hook, "j"] \arrow[drr, bend left, "p'"] \arrow[dd, "q'"'] & & \\
 & G/P \times \mathbb{A}^n \arrow[r, "p"] \arrow[d, "q"'] & G/P \\
\mathrm{Spec}\, R \arrow[r, hook, "i"] & \mathbb{A}^n &
\end{tikzcd}$$

Lemma 2.5. *Assume $p : X \to Y$ is the projection of a vector bundle over an algebraic scheme Y, $\mathscr{T}$ is a quasi-coherent sheaf over Y such that $\mathscr{T}^{\perp} = 0$ in $D^b(Y)$, then,*

$$(p^*\mathscr{T})^{\perp} = 0$$

in $D^b(X)$.

Proof. The projection p is an affine morphism. Denote the corresponding sheaf of algebras of the affine morphism p by $\mathscr{A}$. The push-forward functor p_* induces an equivalence between $Qcoh\mathscr{O}_X$ and the subcategory $\mathscr{A}$*-mod* of $Qcoh\mathscr{O}_Y$. In particular, p_* is exact.

The sheaf $\mathscr{A} = \mathrm{Sym}(X^*)$ is locally free. Therefore, p^* is also exact.

We have a pair of adjoint exact functors (p^*, p_*) between $Qcoh(X)$ and $Qcoh(Y)$. By a standard result in homological algebra (III.6 in [GM03]), they induce an adjoint pair between $D^b(X)$ and $D^b(Y)$, which we still denote by (p^*, p_*). They commute with the shifting functor [1].

For a complex C over X, assume $\mathrm{Hom}_{D^b(X)}(p^*\mathscr{T}, C[i]) = 0$ for all i. Using the adjoint property, we get $\mathrm{Hom}_{D^b(Y)}(\mathscr{T}, p_*C[i]) = \mathrm{Hom}_{D^b(X)}(p^*\mathscr{T}, C[i]) = 0$. By assumption, $p_*C \cong 0$ in $D^b(Y)$. This implies $p_*C \cong 0$ in $D^b(\mathscr{A})$, hence, $C \cong 0$ in $D^b(X)$. ◻

Lemma 2.6. *Assume $p : X \to Y$ is the projection of a vector bundle over an algebraic scheme Y, $\mathscr{T}_1$ and $\mathscr{T}_2$ are two vector bundles over Y such that $H^i(X, p^*\mathscr{H}\mathrm{om}(\mathscr{T}_1, \mathscr{T}_2)) = 0$ for all $i > 0$, then $\mathrm{Ext}^i(p^*\mathscr{T}_1, p^*\mathscr{T}_2) = 0$ for all $i > 0$.*

Proof. Denote the corresponding sheaf of algebras of the affine morphism p by $\mathscr{A}$. We have the local-global spectral sequence E with

$$E_2^{ij} = H^i(X, \mathscr{E}\mathrm{xt}^j_X(p^*\mathscr{T}_1, p^*\mathscr{T}_2))$$

which converges to $\mathrm{Ext}^{i+j}(p^*\mathscr{T}_1, p^*\mathscr{T}_2)$.

Since $p^*\mathscr{T}_1$ and $p^*\mathscr{T}_2$ are both locally free, $\mathscr{E}\mathrm{xt}^j_X(p^*\mathscr{T}_1, p^*\mathscr{T}_2) = 0$ for all positive j. We have

$$\mathrm{Ext}^i(p^*\mathscr{T}_1, p^*\mathscr{T}_2) = H^i(X, \mathscr{H}\mathrm{om}_{\mathscr{O}_X}(p^*\mathscr{T}_1, p^*\mathscr{T}_2)).$$

Since $\mathscr{A}$ is locally free, we can identify

$$p^*\mathscr{H}\mathrm{om}_{\mathscr{O}_Y}(\mathscr{T}_1, \mathscr{T}_2) \cong \mathscr{H}\mathrm{om}_{\mathscr{O}_Y}(\mathscr{T}_1, \mathscr{T}_2 \otimes_{\mathscr{O}_Y} \mathscr{A}) \cong \mathscr{H}\mathrm{om}_{\mathscr{O}_X}(p^*\mathscr{T}_1, p^*\mathscr{T}_2).$$

Use the assumption that $H^i(X, p^*\mathscr{H}\mathrm{om}_{\mathscr{O}_Y}(\mathscr{T}_1, \mathscr{T}_2)) = 0$, then, $\mathrm{Ext}^i(p^*\mathscr{T}_1, p^*\mathscr{T}_2) = 0$ for $i > 0$. ◻

We get the following method to construct crepant noncommutative desingularizations.

Proposition 2.7. *Notations as in diagram (1). Let $\mathscr{T}il$ be a tilting bundle over G/P.*

1. *If $H^i(Z, p'^*\mathscr{E}\mathrm{nd}_{\mathscr{O}_{G/P}}(\mathscr{T}il)) = 0$ for all positive i, then $p'^*\mathscr{T}il$ is a tilting bundle over Z.*
2. *If moreover, $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$ is maximal Cohen-Macaulay, and*

$$\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \mathrm{End}_R(q'_*p'^*\mathscr{T}il),$$

   then $\mathrm{End}_R(q'_*p'^*\mathscr{T}il)$ gives a noncommutative crepant desingularization of $\mathrm{Spec}\, R$.
3. *Assume $H^i(Z, p'^*\mathscr{E}\mathrm{nd}_{\mathscr{O}_{G/P}}(\mathscr{T}il)) = 0$ for all $i > 0$. If the exceptional locus of $q' : Z \to \mathrm{Spec}\, R$ has codimension at least two in both Z and $\mathrm{Spec}\, R$, and R is an integral domain, then $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \mathrm{End}_R(q'_*p'^*\mathscr{T}il)$ and it gives a noncommutative desingularization of $\mathrm{Spec}\, R$.*

Proof. The first part of this proposition follows directly from Lemma 2.5, Lemma 2.6 and the definition of tilting bundles.

Note that if $p'^*\mathscr{T}il$ is a tilting bundle, then $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$ has finite global dimension by Theorem 2.1 because Z is smooth. This shows the second part.

Now let us prove the third part. First note that by Lemma 4.2.1 in [V04a], if the exceptional locus of q' has codimension at least 2 in both Z and $\mathrm{Spec}\, R$, then q'_* sends any reflexive sheaves to reflexive sheaves. Note also that for a finitely generated module M over a commutative noetherian domain, M being reflexive is equivalent to being torsion free and $M = \bigcap M_p$, where the intersection is taken over all codimension 1 primes (7.4.2 of [Bou98]). Both Z and $\mathrm{Spec}\, R$ are integral schemes, hence for any reflexive sheaf N over one or the other, $\mathscr{H}om(M, N)$ is also reflexive for any M. This is because $\mathscr{H}om(M, N)$ is torsion-free whenever N is, and both intersection and localization commute with $\mathscr{H}om(M, -)$, given integrality of the scheme. Thus, we know $p'^*\mathscr{T}il$ is reflexive, so are $\mathscr{E}nd_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$, $q'_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$, $q'_*p'^*\mathscr{T}il$, and $\mathrm{End}_R(q'_*p'^*\mathscr{T}il)$. The map $\mathrm{End}_Z(p'^*\mathscr{T}il) \to \mathrm{End}_S(q'_*p'^*\mathscr{T}il)$ is an isomorphism of rings in the complement of a codimension at least 2 subvariety. Both $\mathrm{End}_Z(p'^*\mathscr{T}il)$ and $\mathrm{End}_S(q'_*p'^*\mathscr{T}il)$ are reflexive, hence $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) = \bigcap \mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)_p \cong \bigcap \mathrm{End}_R(q'_*p'^*\mathscr{T}il)_p = \mathrm{End}_R(q'_*p'^*\mathscr{T}il)$, where the intersection is taken over all codimension 1 primes. □

In some situations we will deal with, the exceptional locus has codimension 1 in Z. Unfortunately Proposition 2.7 doesn't apply. Nevertheless, we have the following Lemma.

Lemma 2.8. *Let $X = \mathrm{Spec}(R)$ be an affine normal Gorenstein scheme and $f : Z \to X$ a crepant resolution with exceptional locus has codimension at least 2 in X. Let $\mathscr{T}il$ be a tilting bundle over Z. If $\mathscr{T}il(Z)$ is reflexive and $\mathrm{End}_{\mathscr{O}_Z}(\mathscr{T}il)$ is maximal Cohen-Macaulay, then*

$$\mathrm{End}_{\mathscr{O}_Z}(\mathscr{T}il) \cong \mathrm{End}_R(\mathscr{T}il(Z))$$

and it is a non-commutative crepant resolution of X.

Proof. The only thing we need to show is that taking global section and taking endomorphism ring commute in this case. First recall that (7.4.2 of [Bou98]) for a finitely generated module M over a commutative noetherian domain, M being reflexive is equivalent to being torsion free and $M = \bigcap M_p$ where the intersection is taken over all codimension 1 primes.

There is a natural morphism of rings $\mathrm{End}_{\mathscr{O}_Z}(\mathscr{T}il) \to \mathrm{End}_R(\mathscr{T}il(Z))$. The target is reflexive since it is a maximal Cohen-Macaulay module over a Gorenstein domain. This morphism is an isomorphism outside the exceptional locus of f which has codimension at least 2 in X. If the source is also reflexive, then we have $\mathrm{End}_{\mathscr{O}_Z}(\mathscr{T}il) = \bigcap \mathrm{End}_{\mathscr{O}_Z}(\mathscr{T}il)_p \cong \bigcap \mathrm{End}_R(\mathscr{T}il(Z))_p = \mathrm{End}_R(\mathscr{T}il(Z))$, where the intersection is taken over all codimension 1 primes.

□

We have the following Lemma to test whether a non-commutative desingularization obtained as in Proposition 2.7 is crepant or not. Although it is known, (e.g., see [BLV11],) we include the proof for completeness.

Lemma 2.9. *Assume Z is the total space of the vector bundle V on G/P, and $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \mathrm{End}_R(q'_*p'^*\mathscr{T}il)$ with $H^i(Z, \mathscr{E}nd_{\mathscr{O}_Z}(p'^*\mathscr{T}il)) = 0$. Then, $\mathrm{End}_R(q'_*p'^*\mathscr{T}il)$ is maximal Cohen-Macaulay iff $H^i(G/P, \mathscr{T}il^* \otimes \mathscr{T}il \otimes \mathrm{Sym}(V^*) \otimes \omega_Z) = 0$, where ω is the dualizing complex.*

Proof. By definition, $\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$ is maximal Cohen-Macaulay iff $\mathrm{Ext}^i_R(\mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il), \omega_R) = 0$ for all $i > 0$. As q is a proper and $Rq_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \mathrm{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$, using the

Grothendieck duality for proper morphisms (1.2.22 of [W03])

$$\mathrm{Ext}^i_R(\mathrm{End}_{\mathcal{O}_Z}(p'^*\mathcal{T}il), \omega_R) \cong \mathrm{Ext}^i_{\mathcal{O}_Z}(\mathcal{E}\mathrm{nd}_{\mathcal{O}_Z}(p'^*\mathcal{T}il), \omega_Z).$$

We know $\mathrm{Ext}^i_R(\mathrm{End}_{\mathcal{O}_Z}(p'^*\mathcal{T}il), \omega_R) = 0$ is equivalent to the following

$$\begin{aligned}\mathrm{Ext}^i_R(\mathrm{End}_{\mathcal{O}_Z}(p'^*\mathcal{T}il), \omega_R) &\cong \mathrm{Ext}^i_{\mathcal{O}_Z}(\mathcal{E}\mathrm{nd}\,\mathcal{O}_Z(p'^*\mathcal{T}il), \omega_Z)\\ &\cong H^i(Z, \mathcal{E}\mathrm{nd}_{\mathcal{O}_Z}(p'^*\mathcal{T}il)\otimes\omega_Z)\\ &\cong H^i(G/P, \mathcal{T}il^*\otimes\mathcal{T}il\otimes\mathrm{Sym}(V^*)\otimes\omega_Z)\\ &\cong 0.\end{aligned}$$

□

2.3. **Exceptional collections and dual collections.**

Definition 2.10 ([HV07]). Let C be a triangulated category. A subset Ω is called spanning if for each object $a \in C$ each of the following conditions implies $a \cong 0$:

1. $\mathrm{Ext}^i(a, b) = 0$ for all $b \in \Omega$ and all $i \in \mathbb{Z}$,
2. $\mathrm{Ext}^i(b, a) = 0$ for all $b \in \Omega$ and all $i \in \mathbb{Z}$.

Let $\mathfrak{A}$ be a k-linear category.

Definition 2.11. An ordered set of objects $\nabla = \{\nabla_\alpha, \alpha \in I\}$ in $D^b(\mathfrak{A})$ is called exceptional if we have $\mathrm{Ext}^\bullet(\nabla_\alpha, \nabla_\beta) = 0$ for $\alpha < \beta$; $\mathrm{Ext}^n(\nabla_\alpha, \nabla_\alpha) = 0$ for $n \neq 0$, and $\mathrm{End}(\nabla_\alpha) = k$. An exceptional set is said to be full if it is spanning for $D^b(\mathfrak{A})$.

Definition 2.12. Let $\Delta = \Delta_\alpha, \alpha \in I$ be another subset of objects in $D^b(\mathfrak{A})$, in bijection with ∇. We say that Δ is the dual collection to ∇ if $\mathrm{Ext}^\bullet(\Delta_\beta, \nabla_\alpha) = 0$ for $\beta > \alpha$, and there exists an isomorphism $\Delta_\beta \cong \nabla_\beta \bmod D_{<\beta}$, where $D_{<\beta}$ is the full triangulated subcategory generated by $\{\nabla_\alpha \mid \alpha < \beta\}$.

There are some well-known facts ([Be06]):

1. $\mathrm{Ext}^\bullet(\Delta_\alpha, \nabla_\alpha) = k$, where k lies in homological degree 0;
2. $\mathrm{Ext}^\bullet(\Delta_\alpha, \nabla_\beta) = 0$ for $\alpha \neq \beta$;
3. The dual collection is unique if it exists at all.

In particular, in the definition above, we take $\mathfrak{A}$ to be the category of coherent sheaves over some scheme. It's not hard to see that if we have a finite full exceptional set $\nabla = \{\nabla_\alpha, \alpha \in I\}$ consisting of vector bundles, then $T = \oplus_\alpha \nabla_\alpha$ is a tilting bundle.

2.4. **Shapes of quivers for the non-commutative desingularization.**

Let G be a reductive group, $P < G$ a parabolic subgroup. Suppose $p' : Z \to G/P$ is a vector bundle. Let $u : G/P \to Z$ be the zero section. Let $\nabla(G/P) = \{\nabla_\alpha | \alpha \in I\}$ be a full exceptional collection over G/P and $\Delta(G/P) = \{\Delta_\alpha \mid \alpha \in I\}$ be the dual collection.

Let us start with a provisional method to get the shape of quivers for the non-commutative desingularizations.

Lemma 2.13. *Let $\mathcal{T}il = \oplus_\alpha \nabla_\alpha$ over G/P. Suppose $p'^*\mathcal{T}il$ is a tilting bundle over Z. Then,* $\mathrm{Ext}^t_{\mathcal{O}_Z}(p'^*\nabla_\alpha, u_*\Delta_\beta)$ *vanishes unless $t = 0$ and $\alpha = \beta$ in which case it is 1-dimensional.*

In particular, $R\,\mathrm{Hom}_{\mathcal{O}_Z}(p'^*\mathcal{T}il, u_*\Delta_\beta) = \mathrm{Hom}_{\mathcal{O}_Z}(p'^*\nabla_\beta, u_*\Delta_\beta) = k$.

Proof. Note that $\mathrm{Ext}^t_{\mathcal{O}_Z}(p'^*\nabla_\alpha, u_*\Delta_\beta) = \mathrm{Ext}^t_{\mathcal{O}_{G/P}}(\nabla_\alpha, \Delta_\beta)$. The conclusion then follows from the facts (1) and (2) about dual exceptional collections listed above. □

From this lemma, we easily get the following Corollary.

Corollary 2.14. *Suppose further more that* $\{R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha), \alpha \in I\}$ *is a complete set of (distinct) indecomposable projective objects in* $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$*-mod.*

Then, for each $\alpha \in I$, $u_*\Delta_\alpha$ *is the simple object corresponds to the indecomposable projective* $R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$.

Remark 2.15. Even if $\nabla(G/P)$ is a full exceptional collection over G/P, the collection of inverse images $\{p'^*\nabla_\alpha \mid \alpha \in I\}$ is rarely an exceptional collection over Z. This is because there could be non-trivial maps $p'^*\nabla_\alpha \to p'^*\nabla_\alpha$ for $\alpha < \beta$.

Throughout this paper, by *radical*, we mean the Jacobson radical.

Proposition 2.16. *Let* $\Lambda := \operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$*. Assume* $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \operatorname{End}_R(q_*p'^*\mathscr{T}il)$ *and* $\operatorname{End}_R(q_*p'^*\mathscr{T}il)$ *is a non-commutative desingularization of* $\operatorname{Spec} R$*. Let* $S_\alpha = R\operatorname{Hom}(p'^*\mathscr{T}il, u_*\Delta_\alpha)$*. Then* S_α*'s are distinct simple objects in* Λ*-mod.*

Assume further that $\Lambda/\operatorname{rad}(\Lambda)$ *is finite dimensional over k and* Λ *is Krull-Schmidt, semi-perfect (projective covers exist in* Λ*-mod). Let* P_α *be the projective cover of* S_α*. Then,* $P_\alpha \cong R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$*, and* $\Lambda = \operatorname{End}_\Lambda(\oplus_\alpha P_\alpha)$ *is a basic algebra.*

See [ASS06] for the definitions of basic algebras.

Note that if $A/\operatorname{rad}(A)$ is finite dimensional, then its radical is zero and hence its module category is semi-simple. We record here some facts that will be used in the proof of the proposition below.

Proposition 2.17. *Notations as above.*

1. *For any A-module M,* $\operatorname{rad} M = \operatorname{rad} A \cdot M$*;*
2. *The assignment* $M \mapsto M/\operatorname{rad} M$ *defines a functor from A-mod to* $A/\operatorname{rad} A$*-mod;*
3. *For any idempotent element* $e \in A$*, the natural map* $\operatorname{Hom}_A(Ae, M) \to Me$ *is an isomorphism of eAe-modules;*
4. *For any* $A/\operatorname{rad} A$*-modules M and N, let* P_N *be the projective cover of N in A-mod, we have* $\operatorname{Hom}_A(N, M) \cong \operatorname{Hom}_A(P_N, M)$*;*
5. *For any* $A/\operatorname{rad} A$*-module M, any simple* $A/\operatorname{rad} A$*-module S and the projective cover P of S in A-mod, we have* $\operatorname{Hom}_A(M, S) \cong \operatorname{Hom}_A(P, M)$.

In the case of finite dimensional algebra, this proposition can be found in [ASS06]. But the same proof applied in this set-up.

Proof of Proposition 2.16. Recall that $R\operatorname{Hom}(p'^*\mathscr{T}il, -)$ induces an equivalence between derived categories. Note that for any α, $R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$ is concentrated in degree 0, and isomorphic to Λe_α, where $e_\alpha \in \Lambda$ is the identity element of $\operatorname{End}_{\mathscr{O}_Z}(p'^*\nabla_\alpha)$. Up to isomorphism, there are no more indecomposable projective Λ-modules because $\Lambda \cong \oplus\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$ and they are direct summands of Λ. Note also that $S_\alpha = R\operatorname{Hom}(p'^*\mathscr{T}il, u_*\Delta_\alpha)$ are all concentrated in degree 0 as well, and each isomorphic to $\mathbb{C}$. Hence all of S_α's are simple objects in Λ-mod. They are pairwise non-isomorphic, because $u_*\Delta_\alpha$'s are distinct, as can be seen from Lemma 2.13.

Note that P_α's form a complete set of (distinct) indecomposable projectives. They are distinct as their tops are distinct. Clearly they are indecomposable as their tops are. These are all the indecomposable projectives (up to isomorphism) since

$$\#I = \operatorname{rank} K_0(G/P) = \operatorname{rank} K_0(Z) = \operatorname{rank} K_0(\Lambda\text{-}mod) = \#\{P_\alpha \mid \alpha \in I\},$$

where the 2nd equality comes from the Thom isomorphism (5.4 of [CG00]) and the 3rd one from the derived equivalence (i.e., any abelian category has the same K-group as its derived category).

It remains to show that $\Lambda e_\alpha \cong P_\alpha$. As a projective module, Λe_α can be decomposed as direct sum of indecomposable projectives. And Lemma 2.13 yields $\Lambda e_\alpha \cong P_\alpha$. □

Remark 2.18. It is folklore that under the assumptions of the second part of Proposition 2.16, in the quiver with relation for $\Lambda = \operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$, the arrows are given by a basis of $\operatorname{Ext}^1_{\mathscr{O}_Z}(\Delta_i, \Delta_j)^*$. More precisely, $e_\alpha(\operatorname{rad}\Lambda/\operatorname{rad}^2\Lambda)e_\beta \cong \operatorname{Ext}^1_\Lambda(S_\alpha, S_\beta)^* \cong \operatorname{Ext}^1_{\mathscr{O}_Z}(\Delta_i, \Delta_j)^*$. And the relations are given by $\operatorname{Ext}^2_{\mathscr{O}_Z}(\Delta_i, \Delta_j)^*$.

Unfortunately, the conditions in Proposition 2.16 and Remark 2.18 are rarely satisfied. We will see later on that there could be infinitely many simple modules over $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$. However, the conclusion of Remark 2.18 is sometimes true in the situations we consider, as we assume that Z is the total space of an equivariant vector bundle V over G/P, and both ∇ and Δ consist of equivariant sheaves over G/P. Note that in this case, $\Lambda = \operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$ is a representation of G.

Proposition 2.19. *Let $\nabla(G/P) = \{\nabla_\alpha \mid \alpha \in I\}$ be a full exceptional collection consisting of equivariant sheaves over G/P with the dual collection $\Delta(G/P)$ also equivariant. Assume the resolution $q : Z \to \operatorname{Spec} R$ is G-equivariant with $q^{-1}(0) = G/P$, and the only G-fixed closed point of $\operatorname{Spec} R$ is $\{0\} \subset \operatorname{Spec} R$. Let $\mathscr{T}il = \oplus_\alpha \nabla_\alpha$. Assume moreover that $p'^*\mathscr{T}il$ is a tilting bundle over Z such that $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il) \cong \operatorname{End}_R(q_*p'^*\mathscr{T}il)$.*

Then, the only G-equivariant simple $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$ modules are given by

$$S_\beta := R\operatorname{Hom}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha), u_*\Delta_\beta) \cong \operatorname{Hom}_{\mathscr{O}_Z}(p'^*\nabla_\beta, u_*\Delta_\beta)$$

and they are pairwise distinct as $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$-modules.

Proof. First we show that the simple equivariant $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$ modules have to be scheme theoretically supported on $i : \{0\} \hookrightarrow \operatorname{Spec} R$. Let M be a simple equivariant $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$ module whose support contains $\{0\}$. Then M is an R-module. The surjective morphism $M \twoheadrightarrow i_*i^*M$ has non-trivial target, because $\{0\}$ is a subset of the support of M. Note that his map is an G-equivariant morphism of $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$-modules. It has trivial kernel since M is simple. Thus, $M \cong i_*i^*M$, i.e., M has (scheme theoretical) support on $\{0\}$.

Now we know that M, with support on $\{0\}$, is a simple module over

$$i^*\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha)) \cong i^*q_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha)).$$

Using the cartesian diagram

$$\begin{tikzcd} Z \arrow[d, "q"'] & G/P \arrow[l, hook', "u"'] \arrow[d, "q'"] \\ \operatorname{Spec} R & \{0\} \arrow[l, hook', "i"] \end{tikzcd}$$

and Remark III.9.3.1 in [H77], we obtain a map

$$i^*q_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha)) \to q'_*u^*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha)).$$

Note that $i^*q_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$, as a inverse image through proper morphisms, is a finite dimensional algebra. There's a natural grading on it giving by the weights of $\mathbb{G}_m$-action, where $\mathbb{G}_m$ acts on Z by scaling. The zero's graded piece of $i^*q_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$ is $\operatorname{End}_{G/P}(\oplus\nabla_\alpha)$. Every element in the first graded piece can be easily checked to be in the Jacobson radical of $i^*q_*\mathscr{E}nd_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$ using finite dimensionality. Therefore, we get an isomorphism

$$i^*\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))/\operatorname{rad} \cong \operatorname{End}_{G/P}(\oplus\nabla_\alpha)/\operatorname{rad}.$$

So far we know M is a simple module over $\operatorname{End}_{G/P}(\oplus\nabla_\alpha)$. As ∇ is a full exceptional collection over G/P, its endomorphism ring is a finite dimensional basic algebra, with all the simple modules of the form $\operatorname{Hom}_{G/P}(\oplus\nabla_\alpha, \Delta_\beta) \cong \operatorname{Hom}_Z(p'^*(\oplus_\alpha\nabla_\alpha), u_*\Delta_\beta)$. Thus, M has to be isomorphic to one of them. (Note that the jacobson radical acts trivially on the simple objects. Thus, two simple modules are isomorphic over $\operatorname{End}_{G/P}(\oplus\nabla_\alpha)$ iff they are isomorphic over $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$.)

For the claim that $S_\alpha \neq S_\beta$ unless $\alpha = \beta$, note that $R\operatorname{Hom}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha), -)$ induces an equivalence of derived categories. To show $S_\alpha \neq S_\beta$, it suffices to show $u_*\Delta_\alpha \neq u_*\Delta_\beta$, and this is clear by Lemma 2.13. □

Remark 2.20. As can be seen from the proof, the finite dimensional algebra $\operatorname{End}_{\mathscr{O}_{G/P}}(\oplus\nabla_\alpha)$ is a subring of $\operatorname{End}_{\mathscr{O}_Z}(p'^*(\oplus_\alpha\nabla_\alpha))$. The modules S_α are also simple modules over $\operatorname{End}_{\mathscr{O}_{G/P}}(\oplus\nabla_\alpha)$. Actually, it is easier to see that S_α's are distinct as modules over $\operatorname{End}_{\mathscr{O}_{G/P}}(\oplus\nabla_\alpha)$.

Remark 2.21. In Proposition 2.19 we only characterized all the simple $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$-modules which happen to be equivariant. There could be more simple objects in the abelian category of G-equivariant $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$-modules.

As $R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$ could be different than the projective covers of $\mathbb{S}_\alpha$, from now on we make the convention that by P_α we mean $R\operatorname{Hom}(p'^*\mathscr{T}il, p'^*\nabla_\alpha)$.

Definition 2.22. For an algebra with a rational G-action, and any G-equivariant module M, we define $\operatorname{rad}_G M$ to be the intersection of all the G-equivariant maximal submodules of M.

Caution: Note that $\operatorname{rad}_G M$ is not the intersection of all the maximal subobjects of M in the category of G-equivariant modules.

Lemma 2.23. *Assume Λ is a k-algebra with a rational G-action such that $\Lambda/\operatorname{rad}_G\Lambda$ is semi-simple. For any equivariant module M over $\Lambda = \operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$, we have $\operatorname{rad}_G M = \operatorname{rad}_G\Lambda\cdot M$.*

Proof. To show the inclusion $\operatorname{rad}_G M \supseteq \operatorname{rad}_G\Lambda\cdot M$, it suffices to show that $\operatorname{rad}_G\Lambda$ is in the kernel of the map $m\cdot : \Lambda \to M/N$ (multiplication by m) for any maximal submodule N and any element $m \in M$. This is true because M/N is simple.

Because of the semi-simplicity of $\Lambda/\operatorname{rad}_G\Lambda$, $M/\operatorname{rad}_G M$ decomposes into direct sum of equivariant simple objects. And it is clear that $\operatorname{rad}_G M$ is the maximal submodule with this property. Hence we get the reverse inclusion. □

Remark 2.24. Up to now, we have not used the fact that the characteristic of the ground field is zero. But in the following theorem we do use.

Theorem 2.25. *Let $\nabla(G/P) = \{\nabla_\alpha \mid \alpha \in I\}$ be a full exceptional collection consisting of equivariant sheaves over G/P with the dual collection $\Delta(G/P)$. Assume the resolution $Z \to \operatorname{Spec} R$ is G-equivariant with $q^{-1}(0) = G/P$, and the only fixed closed point of $\operatorname{Spec} R$ is $\{0\} \subset \operatorname{Spec} R$. Let $\Lambda := \operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$. Assume moreover that $p'^*(\oplus_\alpha\nabla_\alpha)$ is a tilting bundle over Z such that $\Lambda \cong \operatorname{End}_R(q_*p'^*\mathscr{T}il)$. Then,*

1. *we have $e_\alpha(\operatorname{rad}_G\Lambda/\operatorname{rad}_G^2\Lambda)e_\beta \cong \operatorname{Ext}^1_\Lambda(S_\alpha, S_\beta)^*$. In particular, a lifting of a basis of this vector space to $\operatorname{rad}_G\Lambda$ generates Λ over $\oplus_\alpha\mathbb{C}_\alpha$.*
2. *With generators of $\operatorname{End}_{\mathscr{O}_Z}(p'^*\mathscr{T}il)$ chosen as above, $\operatorname{Ext}^2_{\mathscr{O}_Z}(\Delta_i, \Delta_j)^*$ generates the relations.*

Proof of (1). We know that S_α's are all the simple Λ-modules which are G-equivariant. As $\mathrm{Hom}_\Lambda(P_\beta, S_\alpha) = 0$ unless $\alpha = \beta$, in which case $\mathrm{Hom}_\Lambda(P_\beta, S_\alpha) = \mathbb{C}$ and $P_\alpha / \operatorname{rad}_G P_\alpha \cong S_\alpha$. Note that this also implies $\Lambda / \operatorname{rad}_G \Lambda \cong \oplus \mathbb{C}_\alpha$ which is semi-simple.

We take the short exact sequence

$$0 \to \operatorname{rad}_G P_\alpha \to P_\alpha \to S_\alpha \to 0,$$

with $P_\alpha = R\,\mathrm{Hom}(p'^* \mathscr{T}il, p'^* \nabla_\alpha)$. Hence we get an exact sequence

$$0 \longrightarrow \mathrm{Hom}(S_\alpha, S_\beta) \longrightarrow \mathrm{Hom}(P_\alpha, S_\beta) \xrightarrow{0} \mathrm{Hom}(\operatorname{rad}_G P_\alpha, S_\beta) \longrightarrow \mathrm{Ext}^1(S_\alpha, S_\beta) \longrightarrow 0.$$

Hence

$$\begin{aligned} \mathrm{Ext}^1_\Lambda(S_\alpha, S_\beta) &\cong \mathrm{Hom}_\Lambda(\operatorname{rad}_G P_\alpha, S_\beta) \\ &\cong \mathrm{Hom}_\Lambda(\operatorname{rad}_G P_\alpha / \operatorname{rad}^2_G P_\alpha, S_\beta) \\ &\cong \mathrm{Hom}_\Lambda(\Lambda e_\beta, \operatorname{rad}_G P_\alpha / \operatorname{rad}^2_G P_\alpha)^* \\ &\cong \mathrm{Hom}_\Lambda(\Lambda e_\beta, (\operatorname{rad}_G \Lambda / \operatorname{rad}^2_G \Lambda) e_\alpha)^* \\ &\cong e_\alpha (\operatorname{rad}_G \Lambda / \operatorname{rad}^2_G \Lambda) e^*_\beta. \end{aligned}$$

The claim that $\operatorname{rad}_G \Lambda$ generates Λ over $\oplus_\alpha \mathbb{C}_\alpha$ is clear from Lemma 2.23. □

The second part of this theorem follows directly from the next Lemma, which gives an equivariant projective resolution of S_α and is interesting in its own right.

Lemma 2.26. *There is a projective resolution of S_α of the form*

$$0 \longleftarrow S_\alpha \longleftarrow P_\alpha \longleftarrow \oplus_\beta \mathrm{Ext}^1(S_\alpha, S_\beta)^* \otimes P_\beta \longleftarrow \oplus_\beta \mathrm{Ext}^2(S_\alpha, S_\beta)^* \otimes P_\beta \longleftarrow \cdots.$$

Proof. We start with a surjective map $S_\alpha \twoheadleftarrow P_\alpha$ whose kernel is $\operatorname{rad}_G P_\alpha$ which is a direct sum of finite dimensional representations of G. This tells us that there exist a collection of finite dimensional representations $\{V^1_\gamma\}_\gamma$ of G equipped with an equivariant surjective map $\operatorname{rad}_G P_\alpha \twoheadleftarrow \oplus_\gamma V^1_\gamma \otimes P_\gamma$ which fits into an exact sequence

$$0 \leftarrow S_\alpha \leftarrow P_\alpha \leftarrow \oplus_\gamma V^1_\gamma \otimes P_\gamma.$$

We take the kernel and proceed to get an equivariant projective resolution of S_α with i-th term of the form $\oplus_\gamma V^i_\gamma \otimes P_\gamma$.

We apply $\mathrm{Hom}_\Lambda(-, S_\beta)$ to get

$$0 \to 0 \to \delta^\beta_\alpha \mathbb{C} \to V^{1*}_\beta \to V^{2*}_\beta \to \cdots,$$

which is a chain complex of G-representations. Hence, its i-th homology can be identified with a quotient of the i-th term. In particular, we can do this for $i = 1$ and for all β. We obtained

$$\begin{tikzcd}
 & & 0 & \mathrm{coker} \arrow[l] & \\
0 & S_\alpha \arrow[l] & P_\alpha \arrow[l] \arrow[u] & \oplus_\gamma V^1_\gamma \otimes P_\gamma \arrow[l] \arrow[u, two heads, "\pi_1"] & \oplus_\gamma V^2_\gamma \otimes P_\gamma \arrow[l, "d_2"'] \\
 & & & \oplus \mathrm{Ext}^1(S_\alpha, S_\gamma) \otimes P_\gamma \arrow[u, hook] &
\end{tikzcd}$$

We claim that the composition $d_2 \circ \pi_1 : \oplus_\gamma V_\gamma^2 \otimes P_\gamma \to$ coker is surjective. If so, we can replace $V_\gamma^1 \otimes P_\gamma$ by $\mathrm{Ext}^1(S_\alpha, S_\gamma) \otimes P_\gamma$, and $\oplus_\gamma V_\gamma^2 \otimes P_\gamma$ by $d_2^{-1}(\mathrm{Ext}^1(S_\alpha, S_\gamma) \otimes P_\gamma)$, and proceed further to get the desired resolution.

To prove the surjectivity of $d_2 \circ \pi_1$, we can assume coker $= V_\gamma^{1'} \otimes P_\gamma \neq 0$, hence $V_\gamma^{1'} \neq 0$ for some γ. We show that $V_\gamma^{1'} \otimes P_\gamma$ is in the image of $d_2 \circ \pi_1$. Let $W_\gamma := V_\gamma^{1'} \otimes P_\gamma / V_\gamma^{1'} \otimes P_\gamma \cap im(d_2 \circ \pi_1)$. If this isn't zero, $M/\operatorname{rad}_G M$ would contribute to $\mathrm{Ext}^1(S_\alpha, S_\beta)$ which makes it larger than it actually is. Hence, we are done. □

Remark 2.27. Note that the complexes $u_*\Delta_\alpha$'s are not in the core of the usual t-structure in $D^b(Coh(Z))$, but $R\operatorname{Hom}(p'^*\mathscr{T}il, u_*\Delta_\alpha)$'s are for the usual t-structure of $D^b(\Lambda\text{-}mod)$. This means the functor $R\operatorname{Hom}(p'^*\mathscr{T}il, -)$ doesn't restrict to an equivalence $Coh(Z) \to \Lambda\text{-}mod$.

Remark 2.28. This derived equivalence gives the triangulated category $D^b(\Lambda\text{-}mod)$ a t-structure, by lifting the tautological t-structure of $D^b(\Lambda\text{-}mod)$ (see e.g. IV.4 in [GM03] for the definitions of t-structures of a triangulated categories and the tautological t-structures of derived categories). Note that this t-structure is different than the one given in [Be06] which is called the exotic t-structure. One evidence is that the t-structure we give here doesn't have a finite length core, while the abelian category of exotic sheaves does.

Note that twisting the tilting bundle with any line bundle will give the same endomorphism ring. This tells us we will get the same non-commutative desingularization by taking any exceptional collection in the same $H^1(G/P, \mathscr{O}^*)$ orbit. But the induced t-structures on $D^b(Coh(Z))$ do vary.

3. GRASSMANNIANS

3.1. **A geometric technique.** Now we review a geometric technique used throughout this paper. The reference for this subsection is [W03].

The Schur functor for a partition α applied to a vector space E will be denoted by $\mathbb{S}_\alpha E$ [F97].

Let $h : \mathcal{E} \to X$ be a vector bundle, $\mathrm{Grass}(n-r, \mathcal{E}) \to X$ be corresponding grassmannian bundle of $(n-r)$-planes. Let $0 \to \mathcal{R} \to \mathcal{E} \to \mathcal{Q} \to 0$ be the tautological sequence on $\mathrm{Grass}(n-r, \mathcal{E})$.

We will call the elements of $\mathbb{Z}^n$ the GL_n-weights, or simply weights. A weight α is called dominant if it is non-increasing. It is called $(n-r)$-dominant if $\alpha_1 \geq \cdots \geq \alpha_{n-r}$ and $\alpha_{n-r+1} \geq \cdots \geq \alpha_n$. For an $(n-r)$-dominant weight we can consider two weights $\beta = (\alpha_1, \cdots, \alpha_{n-r})$ and $\gamma = (\alpha_{n-r+1}, \cdots, \alpha_n)$. We define the vector bundle $\mathcal{V}(\alpha) = \mathbb{S}_\beta \mathcal{R}^* \otimes \mathbb{S}_\gamma \mathcal{Q}^*$.

The symmetric group Σ_n acts on the set of weights. Let $\alpha = (\alpha_1, \cdots, \alpha_n)$. The permutation $\sigma_i = (i, i+1)$ acts on the set of weights by: $\sigma_i.\alpha = (\alpha_1, \cdots, \alpha_{i-1}, \alpha_{i+1} - 1, \alpha_i + 1, \alpha_i + 2, \cdots, \alpha_n)$.

Let $\alpha \in \mathbb{Z}^n$ be a dominant integral weight. We denote $\overline{\alpha} = (\alpha_1 - \alpha_n, \ldots, \alpha_{n-1} - \alpha_n, 0)$. By definition the weight $\overline{\alpha}$ is a partition.

Theorem 3.1 (Kempf)**.** *Let K be a field of characteristic $p > 0$. Let $h : \mathrm{Grass}(\mathcal{E}) \to X$ be a structure map. Consider the integral dominant weight and the corresponding vector bundle $\mathcal{V}(\alpha)$ on $\mathrm{Grass}(\mathcal{E})$ defined as above. Then $R^0 h_* \mathcal{V}(\alpha) = \mathbb{S}_{\overline{\alpha}} \mathcal{E}^* \otimes (\wedge^n \mathcal{E}^*)^{\otimes \alpha_n}$ and $R^i h_* \mathcal{V}(\alpha) = 0$ for $i > 0$.*

If $char(k) = 0$, we know more about cohomologies of bundles corresponding to non-dominant weights.

Theorem 3.2 (Bott)**.** *We consider the weight α satisfying $\alpha_i \geq \alpha_{i+1}$ for $i \neq n-r$ and the corresponding vector bundle $\mathcal{V}(\alpha)$ over $\mathrm{Grass}(n-r,\mathcal{E})$ defined above. Then one of the two mutually exclusive possibilities occurs:*

1. *There exists a $\sigma \in \Sigma_n, \sigma \neq 1$, such that $\sigma(\alpha) = \alpha$. Then the higher direct images $\mathcal{R}^i h_* \mathcal{V}(\alpha)$ are zero for $i \geq 0$.*
2. *There exists a unique $\sigma \in \Sigma_n$ such that $\sigma(\alpha) := (\beta)$ is a partition (i.e. is nonincreasing). In this case all higher direct images $\mathcal{R}^i h_* \mathcal{V}(\alpha)$ are zero for $i \neq l(\sigma)$, and*
$$\mathcal{R}^{l(\sigma)} h_* \mathcal{V}(\alpha) = \mathbb{S}_\beta \mathcal{E}^*$$

For a projective variety V of dimension m, $X = \mathbb{A}_k^N$ an affine space, the space $X \times V$ can be viewed as the total space of the trivial vector bundle $\mathcal{E}$ of dimension N over V. Let us consider the subvariety Z in $X \times V$ which is the total space of a subbundle $\mathcal{S}$ in $\mathcal{E}$. We denote by q the projection $q : X \times V \to X$ and by q' the restriction of q to Z. Let $Y = q(Z)$. We get the basic diagram

$$\begin{tikzcd}
Z \arrow[r, hook] \arrow[d, "q'"'] & X \times V \arrow[d, "q"'] \\
Y \arrow[r, hook] & X
\end{tikzcd}$$

The projection from $X \times V$ onto V is denoted by p, and the quotient bundle $\mathcal{E}/\mathcal{S}$ by $\mathcal{T}$. Thus we have the exact sequence of vector bundles on V,

$$0 \to \mathcal{S} \to \mathcal{E} \to \mathcal{T} \to 0$$

The coordinate ring of X will be denoted by A. It is a polynomial ring in N variables over k. We will identify the sheaves on X with A-modules. The direct image $p_*(\mathcal{O}_Z)$ can be identified with the sheaf of algebras $\mathrm{Sym}(\eta)$, where $\eta = \mathcal{S}^*$. For a vector bundle $\mathcal{V}$ over V, the $\mathcal{O}_{X\times V}$-module $\mathcal{O}_Z \otimes p^*\mathcal{V}$ will be denoted by $M(\mathcal{V})$.

Theorem 3.3 (5.1.2 in [W03])**.** *For a vector bundle $\mathcal{V}$ on V, we define free graded A-modules*

$$F(\mathcal{V})_i = \bigoplus_{j\geq 0} H^j(V, \bigwedge^{i+j} \xi \otimes \mathcal{V}) \otimes_k A(-i-j)$$

where $\xi = \mathcal{T}^*$ and (i) means shifting by i.

1. *There exist minimal differentials*
$$d_i(\mathcal{V}) := F(\mathcal{V})_i \to F(\mathcal{V})_{i-1}$$
of degree 0 such that $F(\mathcal{V})_\bullet$ is a complex of free graded A-modules with
$$H_{-i}(F(\mathcal{V})_\bullet) = \mathcal{R}^i q_* M(\mathcal{V}).$$
In particular, the complex $F(\mathcal{V})_\bullet$ is exact in positive degrees.
2. *The sheaf $\mathcal{R}^i q_* M(\mathcal{V})$ is equal to $H^i(Z, M(\mathcal{V}))$ and it can be also identified with the graded A-module $H^i(V, \mathrm{Sym}(\eta) \otimes \mathcal{V})$.*
3. *If $\phi : M(\mathcal{V}) \to M(\mathcal{V}')(n)$ is a morphism of graded sheaves, then there exists a morphism of complexes*
$$f_\bullet(\phi) : F(\mathcal{V})_\bullet \to F(\mathcal{V}')_\bullet(n)$$
Its induced map $H_{-i}(f_\bullet(\phi))$ can be identified with the induced map
$$H^i(Z, M(\mathcal{V})) \to H^i(Z, M(\mathcal{V}'))(n).$$

This theorem will be mentioned as the basic theorem of geometric method in this paper.

Now we come to a criterion for maximal Cohen-Macaulayness in the context of geometric technique. The proof, which can be found in 5.1.5 of [W03], is based on the basic theorem of geometric technique and Lemma 2.9.

Proposition 3.4. *Let* $\mathcal{V}$ *be a bundle over* V*, and let* $\check{\mathcal{V}} := \omega_V \otimes \wedge^{top}\xi \otimes \mathcal{V}^*$*. Assume* $\dim Z = \dim Y$ *and* $R^i q'_*(\mathcal{O}_z \otimes p^*\mathcal{V}) = 0$ *for all* $i > 0$*. Then,* $R^0 q'_*(\mathcal{O}_z \otimes p^*\mathcal{V})$ *is a maximal Cohen-Macaulay module supported on* Y *iff* $R^i q'_*(\mathcal{O}_z \otimes p^*\check{\mathcal{V}}) = 0$ *for all* $i > 0$*.*

We will use this proposition in Section 8 to prove that some non-commutative desingularizations we study are crepant.

3.2. **Exceptional collections on Grassmannians.**

Kapranov [K88] constructed an exceptional collection over $\mathrm{Grass}_{n-r}(E)$, the grassmannian of $(n-r)$-planes in the vector space E over $\mathbb{C}$.

Let

$$0 \to \mathcal{R} \to F \times \mathrm{Grass} \to \mathcal{Q} \to 0$$

be the tautological exact sequence over Grass. He proved that $\{\mathbb{S}_\alpha \mathcal{R}^* \mid \alpha \text{ sub-partition of } (r^{n-r})\}$ is an exceptional collection, where (r^{n-r}) means the rectangle with r columns and $n-r$ rows, and $\mathbb{S}_\lambda$ is the Schur functor corresponding to the partition λ. Hence, the vector bundle

$$\mathcal{T}il_K = \oplus_{\alpha \in \{\text{sub-partition of } (r^{n-r})\}} \mathbb{S}_\alpha \mathcal{R}^*$$

is a tilting bundle. And the dual exceptional collection is given by

$$\{\mathbb{S}_{\alpha'} \mathcal{Q}[|\alpha|] \mid \alpha \text{ sub-partitions of } (r^{n-r})\},$$

where for a partition α we write α' for its transpose. And for any weight α, we call $\sum_i \alpha_i$ its area which is denoted by $|\alpha|$.

Observe that $- \otimes \wedge^{top}\mathcal{Q}$ and $- \otimes \wedge^{top}\mathcal{R}$ define a $\mathbb{Z}^2$ action on the triangulated category $D^b(Coh(\mathrm{Grass}_{n-r}(F)))$. This action sends one exceptional collection to another and preserves duality.

Applying the $\mathbb{Z}^2$-action we can see that $\nabla(\mathrm{Grass}) = \{\mathbb{S}_\lambda \mathcal{Q}^* \mid \lambda \subset ((n-r)^r)\}$ over Grass is also a full exceptional collection. As can be checked by definition, the dual collection is given by $\Delta(\mathrm{Grass}) = \mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}[(n-r)r - |\alpha|]$, where for a subpartition α of r^{n-r} we write α^c for its complement in the rectangle r^{n-r}.

Now, let us work in the set-up as in diagram 1 by taking G/P to be the grassmannian. We describes the Ext's between the equivariant simples $S_\beta = R\,\mathrm{Hom}_{\mathcal{O}_Z}(p^*(\oplus_\alpha \nabla_\alpha), u_*\Delta_\beta)$ as in Proposition 2.19. We will use that to get the shape of quivers with relations of the non-commutative desingularization. The proof of the following lemma is on the same line as the corresponding one in [BLV09].

Lemma 3.5. *Let* Z *be the total space of the vector bundle* $\mathbb{S}_\delta \mathcal{Q}^*$ *for some partition* δ *such that the conditions in Proposition 2.19 are satisfied for the exceptional collection* $\nabla(\mathrm{Grass}) = \{\mathbb{S}_\lambda \mathcal{Q}^* \mid \lambda \subset ((n-r)^r)\}$*. Let* S_α*'s be the simples as in Proposition 2.19. Then, the Ext's among them are given by*

$$\mathrm{Ext}^t(S_\alpha, S_\beta) \cong \oplus_s \oplus_{\lambda \in \wedge^s \mathbb{S}_\delta} H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_{\beta'} \mathcal{R}^* \otimes \mathbb{S}_{\alpha'} \mathcal{R}),$$

where $\wedge^s \mathbb{S}_\delta$ *stands for the decomposition of* $\wedge^s \mathbb{S}_\delta \mathbb{C}^{n-r}$ *into irreducible representations counting multiplicity.*

Proof. We have

$$\begin{aligned}
\mathrm{Ext}^t(S_\alpha, S_\beta) &= \mathrm{Ext}^t_{\mathscr{O}_{\mathcal{Z}}}(u_*\mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}[(n-r)r-|\alpha|], u_*\mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\beta^c)'}\mathcal{R}[(n-r)r-|\beta|]) \\
&= \mathrm{Ext}^{t-|\beta|+|\alpha|}_{\mathscr{O}_{\mathcal{Z}}}(u_*\mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}, u_*\mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\beta^c)'}\mathcal{R}) \\
&= \oplus_s H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \wedge^s\mathbb{S}_\delta\mathcal{Q}^* \otimes \mathbb{S}_{(\beta^c)'}\mathcal{R} \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}^*) \\
&= \oplus_s \oplus_{\lambda\in\wedge^s\mathbb{S}_\delta} H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_{(\beta^c)'}\mathcal{R} \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}^*) \\
&= \oplus_s \oplus_{\lambda\in\wedge^s\mathbb{S}_\delta)} H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_{\beta'}\mathcal{R}^* \otimes \mathbb{S}_{\alpha'}\mathcal{R}).
\end{aligned}$$

The only thing need to explain in the above is the third equality. To get that we take the Koszul resolution of $u_*V := u_*\mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}$ as follows

$$\cdots \to \wedge^2\mathbb{S}_\delta\mathcal{Q} \otimes V \otimes \mathscr{O}_{\mathcal{Z}} \to \mathbb{S}_\delta\mathcal{Q} \otimes V \otimes \mathscr{O}_{\mathcal{Z}} \to V \otimes \mathscr{O}_{\mathcal{Z}},$$

apply $\mathscr{H}om_{\mathscr{O}_{\mathcal{Z}}}(-, u_*U)$, and use adjunction formula to get

$$0 \to \mathscr{H}om_{\mathrm{Grass}}(V, U) \to \mathscr{H}om_{\mathrm{Grass}}(\mathbb{S}_\delta\mathcal{Q} \otimes V, U) \to \mathscr{H}om_{\mathrm{Grass}}(\wedge^2\mathbb{S}_\delta\mathcal{Q} \otimes V, U) \to \cdots.$$

Note that the t-th hypercohomology of this complex is exactly $\mathrm{Ext}^t(S_\alpha, S_\beta)$. The third equality above is equivalent to the degeneracy of the hypercohomology spectral sequence

$$E_1^{r,s} = H^r(\mathscr{H}om_{\mathrm{Grass}}(\wedge^s\mathbb{S}_\delta\mathcal{Q} \otimes V, U)) \Rightarrow \mathrm{Ext}^{r+s}(S_\alpha, S_\beta)$$

at E_1-page. We calaim that the hypercohomology spectral sequence does degenerate at E_1-page. In fact, after pluging in $V = \mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\alpha^c)'}\mathcal{R}$ and $U = \mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{(\beta^c)'}\mathcal{R}$, all the differentials are equivariant under GL_n and the areas of the weights of different columns in the spectral sequence are different. Therefore, no differentials other than the vertical ones which are in E_0. ◻

Corollary 3.6. *Let $\mathbb{S}_\gamma\mathcal{Q}^* \subset \mathbb{S}_{\beta'}\mathcal{Q}^* \otimes \mathbb{S}_{\alpha'}\mathcal{Q}$ and $k = t - s - |\gamma|$.*

1. *For a fixed t, $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*) = 0$ for any $s > t$.*
2. *For a fixed t, $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*) = 0$ for any $|\delta|s \geq k$.*
3. *For a fixed s and t, $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*) = 0$ for any $\gamma = (\gamma_1, \cdots, \gamma_{n-r})$ with the positive area of γ greater than $t - s$.*
4. *For any $\gamma = (\gamma_1, \cdots, \gamma_{n-r})$, $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*) = 0$ unless the negative part of γ' is contained in λ.*

Proof. According to Bott's Theorem, there can be no more than one k such that $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*) \neq 0$, and this k is the number of adjacent transpositions for $(\gamma_1, \cdots, \gamma_{n-r}, \lambda_1, \cdots, \lambda_r)$ to make it dominant. Assume $H^k \neq 0$, we know $\gamma_{n-r} \geq -r$, and therefore the total negative area in $(\gamma_1, \cdots, \gamma_{n-r})$ is no larger than k. So, $-s + t = |\gamma| + k \geq 0$ which proves (1).

Again because $\gamma_{n-r} \geq -r$, the total area of $(\gamma_1, \cdots, \gamma_{n-r}, \lambda_1, \cdots, \lambda_r)$ is positive. Therefore, $|\delta|s \geq k$.

We know that the total negative area in $(\gamma_1, \cdots, \gamma_{n-r})$ is no larger than k. Let the positive area of γ be l. Then we have $-(|\gamma| - l) \leq k$. Hence $l \leq t - s$.

The last part is clear. ◻

The following examples, which are direct consequences of Corollary 3.6, will be used in Section 5 and Section 6. To state them, we introduce the following notations. For any Young diagram α, by $\tau_l\alpha$ we mean α delating the first l columns, and by τ^l we mean α delating everything after the first l columns. For any partition $\alpha = (\alpha_1, \cdots, \alpha_l)$, by $(-\alpha)$ we mean a partition with $(-\alpha)_i = -\alpha_{l-i}$.

Example 3.7. Notations as above. Let $t = 1$, then, the only non-zero $H^k(\mathrm{Grass}, \mathbb{S}_\lambda\mathcal{Q}^* \otimes \mathbb{S}_\gamma\mathcal{R}^*)$ happens only in the following cases:

- $s = 1$: $-\gamma' = \lambda$ in which case $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{C}$. Or *length*$(\gamma) = n - r$, and $(\tau^{n-r}\lambda) = (-\gamma)'$. In this case, the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{\tau_{n-r}\lambda} E^*$.
- $s = 0$: $\gamma = (1, 0, \cdots, 0)$. In this case, the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = E^*$.

Proof. According to Corollary 3.6, we only need to consider the $s = 1$ and $s = 0$ case. If $s = 1$, then part 3 and 4 Corollary 3.6 yield that $-\gamma' \subset \lambda$. We also have $k = -|\gamma|$. Note that $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) \neq 0$ implies that the weight becomes dominant after k exchanges. Therefore, $(\lambda')_i = (-\gamma)_i$ for all $i \leq$ *length*(γ), and the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{(\lambda_1-(\gamma')_{n-r}, \lambda_2-(\gamma')_{n-r-1}, \cdots, \lambda_{n-r}-(\gamma')_1)} E^*$. The statement for $s = 0$ is clear. □

Similarly, one has the following example.

Example 3.8. Notations as above. Let $t = 2$, then, the only non-zero $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*)$ happens only in the following cases.

- $s = 2$: $-\gamma' = \lambda$ in which case $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{C}$. Or *length*$(\gamma) = n - r$, and $(\tau^{n-r}\lambda) = (-\gamma)'$. In this case, the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{\tau_{n-r}\lambda} E^*$.
- $s = 1$, the positive part of γ is $(1, 0, \cdots, 0)$: The negative part of γ is $(-\lambda)$ in which case $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{C}$. Or the length of the negative part of (γ) is $n - r - 1$, $(\tau^{n-r-1}\lambda) = ((-\gamma')_1, \cdots, (-\gamma')_{n-r-1})$, and $\tau_{n-r-1}\lambda_1 \leq 1$, in which the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{(1, \tau_{n-r-1}\lambda)} E^*$.
- $s = 1$, γ has no positive part: *length*$(\gamma) = n - r$, and $((\tau^{n-r}\lambda)')_i = (-\gamma)_i$ for all $i \leq n-r-1$, $((\tau^{n-r}\lambda)')_{n-r} = (-\gamma)_{n-r} + 1$, and $\lambda_{-\gamma_1+1} - (-\gamma')_{-\gamma_1+1} \geq 2$, in which case the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{(\tau_{n-r}\lambda_1, \cdots, \tau_{n-r}\lambda_{-\gamma_1}, \tau_{n-r}\lambda_{-\gamma_1}, 1, 0, \cdots, 0)} E^*$. Or $\gamma = (0, -1, \cdots, -1)$, $\lambda_2 = 0$ and $\lambda_1 \geq n - r + 1$, in which case the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_{(\lambda_1-n+r, 1, 0, \cdots, 0)} E^*$.
- $s = 0$: $\gamma = (1, 1, 0, \cdots, 0)$ in which case, the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \wedge^2 E^*$. Or $\gamma = (2, 0, \cdots, 0)$ in which case, the corresponding $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) = \mathbb{S}_2 E^*$.

The following lemma and remark gives an algorithm to calculate the higher Ext's. We will fix λ with $|\lambda| = |\delta|s$, and let $t = k - s - |\gamma|$.

Lemma 3.9. (1) *For a fixed partition λ with $l(\lambda) \leq r$, there is a unique dominant GL_{n-r}-weight γ with $\gamma_{n-r} \geq -r$ such that $|\gamma|$ is minimal and $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) \neq 0$ for some k.*

(2) *Let the minimal γ in(1) be $\gamma_{min\lambda}$ and the corresponding t be $t_{min\lambda}$. Every other γ with $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) \neq 0$ for some k has $\gamma_k \geq (\gamma_{min\lambda})_k$ and the corresponding t strictly greater then $t_{min\lambda}$.*

The proof of this lemma shows how to find this minimal γ, the corresponding k and t.

Proof. We look at $\lambda + (r, r-1, \cdots, 1)$. Let $i_0 = r$, for $j > 0$ let $i_j = \lambda_r + r - j - \lambda_{r-j}$, i.e., $\lambda_{r-j} + j + 1 = \lambda_r + 1 + (r - i_j)$. Suppose the largest j with positive i_j is p with the corresponding $i_p = q$. We construct the minimal γ as follows.

Start with $k = r - 1$ and $l = n - r$. If $k = i_j$ for some $j = 1, \cdots, p$, we do nothing for l and decrease k by 1. Otherwise we have $k \neq i_j$ for any $j = 1, \cdots, p$, in which case we set $\gamma_l = \lambda_r + (1 + r - k) - (n - l + 1)$ and then decrease both k and l by 1. Repeat this process. Stop if we reach $l = 0$ or $k = 0$.

If the above process stopped with $k = 0$ and $l \neq 0$, we reset $k = p - 1$ and maintain the same l. If $\lambda_k + (1 + r - k) = \gamma_{l-1} + (n - l + 1) + 1$, we keep the same l and decrease k by 1. Otherwise we have $\lambda_k + (1 + r - k) > \gamma_{l+1} + (n - l + 1) + 1$, in which case we set $\gamma_l = \lambda_k + (1 + r - k) - (n - l + 1) + 1$ and then decrease both k and l by 1. Repeat this process. Stop if we reach $l = 0$ or $k = 0$.

Again if the above process stopped with $k = 0$ and $l \neq 0$, we set $\gamma_l = \gamma_{l+1}$ and decrease l by 1. Repeat this process. Stop if we reach $l = 0$.

The number k is the length of the permutation making (γ, λ) dominant.

The second part is clear from the construction. $\square$

The following remark gives a recursive method to find out all the other γ's with some non-vanishing cohomology.

Remark 3.10. Let the minimal γ in(1) be $\gamma_{min\lambda}$ and the corresponding t be $t_{min\lambda}$. Every other γ with $H^k(\text{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_\gamma \mathcal{R}^*) \neq 0$ for some k can be obtained by a sequence of the following operations $\widehat{\gamma} \mapsto \widehat{\gamma}$.

The operation depends on the parameter $s = 1, \cdots, n - r$. Set $\widehat{\gamma}_t = (\gamma_{min\lambda})_t$ for all $t > s$. Start with $l = s$. We find the largest j with $\max\{\widehat{\gamma}_{l+1} + (n-l) + 1, (\gamma_{min\lambda})_l + (n-l+1) + 1\} < \lambda_j + (r-j+1)$ and set $\widehat{\gamma}_l = \max\{\lambda_{j+1} + (r-j) - (n-l+1) + 1, \widehat{\gamma}_{l+1} + (n-l) + 1, (\gamma_{min\lambda})_l + (n-l+1) + 1\}$. Then we decrease l by 1 and repeat this process. Stop if we reach $l = 0$.

Note that in the calculation of Ext's we only care about those γ's with $\gamma_1 \leq r$. This means we always get all the possible such γ's after finitely many operations above.

4. Equivariant quivers

Here we introduce the basic notions on equivariant quivers and representations of them. They will provide convenient language for the description of non-commutative desingularizations, especially if we would like to consider the equivariant derived categories. Also, as we will see in Subsection 4.2, the derived category of coherent sheaves over homogeneous spaces are easier to describe in this way.

4.1. **The notion of equivariant quivers.** Let G be a reductive group, which will be $GL_n(\mathbb{C})$ later on.

Definition 4.1. An *equivariant quiver* is a triple $Q = (Q_0, Q_1, \alpha)$ where (Q_0, Q_1) is a quiver and α is an assignment associating each arrow $q \in Q_1$ a finite dimensional irreducible representation of G.

Definition 4.2. Let Q be an equivariant quiver. The *path algebra* KQ of Q is the K-algebra whose underlying vector space is

$$\bigoplus_{(q_1,\ldots,q_l) \text{ is a path}} \alpha((q_1, \ldots, q_l)),$$

where $\alpha((q_1, \ldots, q_l)) := \bigotimes_{i=1}^{l} \alpha(q_i)$. We define the product $KQ \otimes KQ \mapsto KQ$ to be

$$\alpha((q_1, \ldots, q_l)) \otimes \alpha((p_1, \ldots, p_h)) \to \alpha((r_1 \ldots, r_s))$$

is given by

$$\begin{cases} \text{id}, \text{ if the path } (r_1 \ldots, r_s) = (q_1, \ldots, q_l, p_1, \ldots, p_h) \\ 0, \text{ otherwise} \end{cases}$$

Obviously, KQ has an action by G and the multiplication is G-equivariant.

We will say Q is finite if both Q_0 and Q_1 are finite sets. We will concentrate on connected finite quiver.

Let Q be a finite quiver, and I be a two sided ideal of KQ. We say (Q, I) is an *equivariant quiver with relations* if I is generated by sub-representations of $\alpha(q_1, \ldots, q_r)$ for paths $(q_1, \ldots, q_r)$. We usually specify a set of such sub-representations as generators of the ideal and call this set relations. We say I is *admissible* if it is generated by sub-representations of $\alpha(q_1, \ldots, q_r)$ for paths of length 2 or longer and I contains some power of the arrow ideal. In this case, the pair (Q, I) will be called a bound equivariant quiver, and KQ/I the bound path algebra.

We would like to define two notions of representations of a bound equivariant quiver, depending on whether or not we will acknowledge G-action.

The first notion will be called simply a representation which is just a representation of the quiver obtained by replacing each arrow $q \in Q_1$ by as many arrows as the dimension of $\alpha(q)$.

Definition 4.3. A *representation* of (Q_0, Q_1, α) is an assignment associating to each vertex $a \in Q_0$ a vector space V_a, and to each arrow $q : a \to b$ a K-linear morphism $V_a \otimes \alpha(q) \to V_b$ (or equivalently $\dim(\alpha(q))$ linear maps $V_a \to V_b$ according to a fixed basis of $\alpha(q)$).

We can as well define the notion of equivariant representations of an equivariant quiver.

Definition 4.4. Let $Q = (Q_0, Q_1)$ be an equivariant quiver. An *equivariant representation* of Q is an assignment associating to each vertex $a \in Q_0$ a representation V_a of G, and to each arrow $q : a \to b$ a G-morphism $V_a \otimes \alpha(q) \to V_b$.

Let (Q, I) be an equivariant quiver with relations. For a bound representation (resp. bound equivariant representation) we require that the generators of I, which we can chose to be sub-representations by definition, act trivially, i.e., all the morphisms above involving subspaces of I are trivial maps.

If A is a commutative K-algebra with a rational G-action, we can also talk about bound representations (resp. bound equivariant representation) of (Q, I) over A. By this we mean associating to each vertex a projective A-module and all the maps have to be A-linear. For equivariant representations we require that the projective modules have a rational G-action compatible with the A-module structure and that all maps corresponding to arrows are G-equivariant.

Proposition 4.5. *The category of bound representations (resp. bound equivariant representations) of (Q, I) (over K) is equivalent to the category of modules (resp. G-equivariant modules) over the ring KQ/I.*

By an equivariant module over KQ/I, we mean an G-equivariant KQ/I action $KQ/I \otimes M \to M$.

4.2. **Beilinson and Kapranov quivers.**

According to a result of Beilinson, we have a full exceptional collection

$$\nabla(\mathbb{P}^{n-1}) := \{\Omega^{i-1}(i) \mid i \in [0, n-1]\}$$

in the derived category of quasi-coherent sheaves over $\mathbb{P}^{n-1}$. Thus, the endomorphism ring of $\oplus_{1=0}^{n-1}\Omega^{i-1}(i)$ is derived equivalent to $\mathbb{P}^{n-1}$, where Ω^k is the k-th exterior power of the sheaf of Kähler differentials.

Let E be a vector space of dimension n and we take $G = GL_n(\mathbb{C})$ acting naturally on E. The Beilinson equivariant quiver, which will be denoted by $QB(n)$, is defined as follows:

$$\begin{tikzcd}
\bullet_0 \arrow[r, bend left, "\alpha_0(E)"] & \bullet_1 \arrow[r, bend left, "\alpha_1(E)"] & \cdots \arrow[r, bend left, "\alpha_{n-1}(E)"] & \bullet_{n-1}
\end{tikzcd}$$

with relations:

$$\alpha_i\alpha_{i+1}(\wedge^2 E).$$

Remark 4.6. Let's pick up a basis for E, say, $e_1, \cdots, e_n$. Then, the above quiver, with equivariant structure forgotten, has n arrows going from the i-th vertex to the $i+1$-th, denoted by $\alpha_i^1, \cdots, \alpha_i^n$, corresponding to the basis elements of E. The relations $\alpha_i\alpha_{i+1}(\wedge^2 E)$ can be written as $\alpha_{i+1}^j\alpha_i^k - \alpha_{i+1}^k\alpha_i^j$ for all j, k.

The bound path algebra of this quiver is isomorphic to $\mathrm{End}_{\mathscr{O}_{\mathbb{P}^{n-1}}}(\oplus_{1=0}^{n-1}\Omega^{i-1}(i))$, as can be checked by $\mathrm{Hom}_{\mathscr{O}_{\mathbb{P}(E)}}(\Omega^{a-1}(a), \Omega^{b-1}(b)) \cong \wedge^{a-b}(E^*)$ if $a \leq b$ and 0 otherwise.

Remark 4.7. It is not hard to see and is proved in [BLV09] that the category of representations of the Beilinson quiver is isomorphic to the category of graded modules over the exterior algebra. Also the same is true for equivariant representations.

The following proposition can be proved using induction.

Proposition 4.8. *The simple equivariant modules over the bound Beilinson quiver $QB(n)$ are of the*

- $V_0 = \mathbb{S}_{\lambda_0}E$ *for some* $\lambda_0 \vdash n$,
- $V_1 = \mathbb{S}_{\lambda_1}E$ *for some* $\lambda_1 \vdash n$ *and* $C^{\lambda_1}_{\lambda_0,(1,0,\cdots,0)} \neq 0$,
- $V_i = \mathbb{S}_{\lambda_i}E$ *for some* $\lambda_i \vdash n$ *and* $C^{\lambda_i}_{\lambda_{i-1},(1,0,\cdots,0)} \neq 0$ *and* $C^{\lambda_i}_{\lambda_{i-2},(1,1,0,\cdots,0)} \neq 0$ *for* $i \geq 2$.

We would like to do the same thing for the grassmannian $\mathrm{Grass}_r(n)$ with $n - r > 1$ and call the corresponding equivariant quiver the Kapranov quiver $QK(r, n)$. The exceptional collection we take will be

$$\nabla(\mathrm{Grass}_r(n)) := \{\mathbb{S}_\alpha\mathcal{R}^*, \alpha \subset ((m-r)^r)\}.$$

Let $G = GL_n(\mathbb{C})$ and $E = \mathbb{C}^n$ with the natural G action.

The Kapranov quiver has the set of vertices corresponding to the set of subpartitions of $((m-r)^r)$, and two vertices λ_1 and λ_2 are linked by an arrow $\lambda_1 \to \lambda_2$ iff λ_2/λ_1 is a single box, and in this case this arrow will be associated to the G-representation E.

For any three partitions α, β, μ, their Littlewood-Richardson coefficient will be denoted by $C^\mu_{\alpha,\beta}$, i.e., the number of Littlewood-Richardson tableaux of shape μ/α and of weight β is $C^\mu_{\alpha,\beta}$. With this notation, the relations in Kapranov quiver are generated by the subrepresentations

$$C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)}\mathbb{S}_2E^* \oplus C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E$$

of the arrows in $\mathrm{Hom}(\beta, \alpha)$.

The bound path algebra of $QK(r, n)$ is isomorphic to $\mathrm{End}_{\mathrm{Grass}}(\oplus_{\alpha\subset((m-r)^r)}\mathbb{S}_\alpha\mathcal{R}^*)$.

Example 4.9. We compute the Kapranov equivariant quiver of the grassmannian $\mathrm{Grass}_2(4)$.

$$\begin{tikzcd}
 & & \ydiagram{1,1} \arrow[dl, "\alpha_2(E)"] & & \\
\varnothing & \ydiagram{1} \arrow[l, "\alpha_1(E)"'] & & \ydiagram{2,1} \arrow[ul, "\alpha_4(E)"'] \arrow[dl, "\alpha_5(E)"] & \ydiagram{2,2} \arrow[l, "\alpha_6(E)"'] \\
 & & \ydiagram{2} \arrow[ul, "\alpha_3(E)"'] & &
\end{tikzcd}$$

with relations:

- Hom($\ydiagram{2}$, $\varnothing$): $\alpha_1\alpha_3(\wedge^2 E)$;
- Hom($\ydiagram{1,1}$, $\varnothing$): $\alpha_1\alpha_2(\mathbb{S}_2 E)$;
- Hom($\ydiagram{2,2}$, $\ydiagram{2}$): $\alpha_5\alpha_6(\wedge^2 E)$;
- Hom($\ydiagram{2,2}$, $\ydiagram{1,1}$): $\alpha_4\alpha_6(\mathbb{S}_2 E)$;
- Hom($\ydiagram{2,1}$, $\ydiagram{1}$): $\alpha_2\alpha_4 - \alpha_3\alpha_5(\mathbb{S}_2 E \oplus \wedge^2 E)$.

Proposition 4.10. *The functor* $\Phi := R\,\mathrm{Hom}_{\mathcal{O}_{\mathrm{Grass}_r(n)}}(\oplus_{\alpha\subset((m-r)^r)}\mathbb{S}_\alpha\mathcal{R}^*, -)$ *induces an equivalence between* $D^b_G(Coh(X))$ *and* $D^b_G(\mathbb{C}QK(r,n)/I\text{-}mod)$*, with quasi-inverse given by* $\Psi := -\otimes^L(\oplus_{\alpha\subset((m-r)^r)}\mathbb{S}_\alpha\mathcal{R}^*)$.

Proof. First note that both functors are well-defined on this level. The Hom space between any two equivariant sheaves is naturally a representation of G and the G-action is compatible with multiplication by elements in $\mathrm{End}_{\mathrm{Grass}}(\oplus_{\alpha\subset((m-r)^r)}\mathbb{S}_\alpha\mathcal{R}^*) \cong \mathbb{C}QK(r,n)/I$, and similar for Ψ. Also both functors commute with the forgetful functors forgetting the G-action. Without the G-equavariance, this result has been discussed. The compatibility of G-actions can be checked directly.

□

5. Determinantal varieties of symmetric matrices

In this section we study a non-commutative desingularization of determinantal varieties in the space of symmetric matrices.

5.1. **Review of the commutative desingularization.**

Now we describe a desingularization of determinantal varieties in the space of symmetric matrices.

Let E be a vector space over k of dimension n and H^s be the subspace of $\mathrm{Hom}_k(E, E^*)$ consisting of symmetric morphisms. This space can be identified with $Sym_2(E^*)$. Upon choosing a set of basis, H^s can be identified with the set of symmetric $(n\times n)$-matrices (x_{ij}) with $x_{ij} = x_{ji}$ the coordinate ring of which can be identified with $S^s = k[x_{ij}]_{i\leq j}$.

Similar to the case of determinantal varieties, we get a universal morphism $\varphi : \mathcal{E} \to \mathcal{E}^*$ over H^s. For $r \leq n$, we desingularize the locus $\mathrm{Spec}\, R^s$, where φ has rank $\leq r$. In other words, R^s is the quotient of S^s by the ideal generated by the $(r+1)\times(r+1)$ minors of (x_{ij}).

Let

$$0 \to \mathcal{R} \to E \times \mathrm{Grass} \to \mathcal{Q} \to 0$$

be the tautological sequence over Grass, where Grass is the grassmannian of $n-r$ planes in E.

With a little abuse of notation, we will add an upper script s for the varieties and keep the same notations for the maps. More precisely, let $\mathcal{Y}^s$ be the product $\mathrm{Grass} \times H^s$, p and q be the projection to Grass and H^s respectively. Inside of $\mathcal{Y}^s$, there is an incidence variety, denoted by $\mathcal{Z}^s$, defined as $\mathcal{Z}^s = \{(g,h) \in \mathrm{Grass} \times H^s : \mathrm{im}\, h \in g\}$. The inclusion $\mathcal{Z}^s \hookrightarrow \mathcal{Y}^s$ is denoted by j, and $\mathrm{Spec}\, R^s \to H^s$ by i. The induced map $\mathcal{Z}^s \to \mathrm{Spec}\, R^s$ from $q : \mathcal{Y}^s \to H^s$ is denoted by q' and $\mathcal{Z}^s \to \mathrm{Grass}$ from $p : \mathcal{Y}^s \to \mathrm{Grass}$ by p'.

These notations are summarized by the following diagram.

$$\begin{tikzcd}
\mathcal{Z}^s \arrow[drr, bend left, "p'"] \arrow[dr, hook, "j"] \arrow[dd, "q'"'] & & \\
& \mathcal{Y}^s = \mathrm{Grass} \times H^s \arrow[r, "p"] \arrow[d, "q"'] & \mathrm{Grass} = \mathrm{Grass}_{n-r}(E) \\
\mathrm{Spec}\, R^s \arrow[r, hook, "i"] & H^s = \mathrm{Sym}_2(E^*) &
\end{tikzcd}$$

Proposition 5.1 (6.3.2 in [W03])**.** *The variety* $\mathcal{Z}^s$ *is a desingularization of* $\mathrm{Spec}\, R^s$.

The variety $\mathcal{Z}^s$ can be described as the total space of the vector bundle $\mathrm{Sym}_2(\mathcal{Q}^*)$ over Grass. Equivalently, $p' : \mathcal{Z}^s \to \mathrm{Grass}$ is an affine morphism with $p_*\mathcal{O}_{\mathcal{Z}^s}$ equal to the sheaf of algebra $\mathrm{Sym}(\mathrm{Sym}_2 \mathcal{Q})$.

5.2. **A tilting bundle over the desingularization.** We consider the inverse image of $\mathcal{T}il_K = \oplus_{\alpha \subset ((n-r)^r)} \mathbb{S}_\alpha \mathcal{Q}^*$, the Kapranov's tilting bundle over $\mathrm{Grass}_{n-r}(E)$, by $p' : \mathcal{Z}^s \to \mathrm{Grass}$. We can show the following.

Lemma 5.2. *For all* $i > 0$, *and any* α, β *sub-partitions of* $((n-r)^r)$, *we have*

$$H^i(\mathcal{Z}^s, p'^* \mathcal{H}om_{\mathcal{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*)) = 0.$$

Proof. Use the Cauchy-Littlewood and the Littlewod-Richardson formulas, we can reduce the sheaf $p'^* \mathcal{H}om_{\mathcal{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*) \cong \mathcal{H}om_{\mathcal{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^* \otimes_{\mathcal{O}_{\mathrm{Grass}}} \mathrm{Sym}(\mathrm{Sym}_2 \mathcal{Q}))$ into the form

$$\oplus (\mathbb{S}_\gamma \mathcal{Q}^*)^{\oplus C_\gamma}$$

for some multiplicities C_γ. By the Bott's theorem 3.2, all the higher cohomology of

$$\mathcal{H}om_{\mathcal{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^* \otimes_{\mathcal{O}_{\mathrm{Grass}}} \mathrm{Sym}(\mathrm{Sym}_2 \mathcal{Q}))$$

vanishes. □

From the lemma above, using Theorem 2.7, we get the following.

Proposition 5.3. *The Kapranov's tilting bundle over* $\mathrm{Grass}_{n-r}(E)$ *is denoted by* $\mathcal{T}il_K$. *The rank* r *determinantal variety* $\mathrm{Spec}\, R^s$ *of symmetric matrices and its desingularization* $\mathcal{Z}^s$ *are as above. The bundle* $p'^* \mathcal{T}il_K$ *is a tilting bundle over* $\mathcal{Z}^s$.

By Theorem 7.6 in [HV07], since $\mathcal{Z}^s$ is smooth, $\mathrm{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K)$ has finite global dimension.

Lemma 5.4. *Notations as above, we have the following.*

1. *If* $r = n-1$, $\mathrm{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K)$ *is maximal Cohen-Macaulay over* $\mathrm{Spec}\, R^s$.
2. *If* $r < n-1$, $\mathrm{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K)$ *is never maximal Cohen-Macaulay over* $\mathrm{Spec}\, R^s$.

Proof. According to Lemma 2.9, it suffices to compute $H^i(\text{Grass}, \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^s} \otimes \text{Sym}(\mathbb{S}_2 \mathcal{Q}))$ for $i > 1$ and $\alpha, \beta \subset (n-r)^r$. The sheaf $\omega_{\mathcal{Z}^s}$ has been computed in 6.7 of [W03], which says $\omega_{\mathcal{Z}^s} \cong \mathbb{S}_{(-n-1+r)^n} E^* \otimes \mathbb{S}_{(n-r-1)^r} \mathcal{Q}^*$. Hence,

$$\mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^s} \otimes \text{Sym}(\mathbb{S}_2 \mathcal{Q}) \cong \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \mathbb{S}_{(-n-1+r)^n} E^* \otimes \mathbb{S}_{(n-r-1)^r} \mathcal{Q}^* \otimes \text{Sym}(\mathbb{S}_2 \mathcal{Q}).$$

Using Cauchy-Littlewood and Littlewood-Richardson formulas, as well as Bott's Theorem, We see that

$$H^i(\text{Grass}, \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^s} \otimes \text{Sym}(\mathbb{S}_2 \mathcal{Q})) = 0$$

if and only if $\mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \cong \oplus \mathbb{S}_\lambda \mathcal{Q}^*$ has $\lambda_1 < 2$ for all λ. This happens for all $\alpha, \beta \subset (n-r)^r$ precisely when $r = n-1$. □

Using the same computation as in Lemma 2.9 and Lemma 5.4, we get

Lemma 5.5. *The R^s-module $q'_* p'^* \mathcal{T}il_K$ is maximal Cohen-Macaulay. In particular, it is reflexive.*

According to Proposition 2.7 and Lemma 2.8, Lemma 5.4 and Lemma 5.5 imply the following Proposition.

Proposition 5.6. *Notations as above, we have the following.*

(1) *The natural map*

$$\text{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K) \to \text{End}_S(q'_* p'^* \mathcal{T}il_K)$$

is an isomorphism of rings.

(2) *If $r = n-1$, $\text{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K)$ is a non-commutative crepant desingularization of $\text{Spec}\, R^s$. If $r < n-1$, $\text{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K)$ is a non-commutative desingularization of $\text{Spec}\, R^s$ (but never crepant).*

Before we show some properties and formulas for this non-commutative desingularization, let us remark about the equivariant projectives and simples over this non-commutative desingularization.

Remark 5.7. In this case, the tilting bundle over $\mathcal{Z}^s$ is the inverse image of an exceptional collection $\nabla(\text{Grass}) = \{\mathbb{S}_\lambda \mathcal{Q}^* \mid \lambda \subset ((n-r)^r)\}$ over Grass. The dual collection is given by $\Delta(\text{Grass}) = \mathbb{S}_{(n-r)^r} \mathcal{Q}^* \otimes \mathbb{S}_{\alpha^{ct}} \mathcal{R}[(n-r)r - |\alpha|]$.

Remark 5.8. Although $p'^*(\mathbb{S}_\alpha \mathcal{Q}^*)$ seems tend to have the same global sections for different α's, they differ as graded objects. This difference is essential as can be seen that if $\alpha \neq \beta$ and $q'_* p'^*(\mathbb{S}_\alpha \mathcal{Q}^*) \cong q'_* p'^*(\mathbb{S}_\beta \mathcal{Q}^*)$, we would have $R\,\text{Hom}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K, p'^*(\mathbb{S}_\alpha \mathcal{Q}^*)) \cong R\,\text{Hom}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K, p'^*(\mathbb{S}_\beta \mathcal{Q}^*))$, i.e., they corresponding to the same object in $D^b(\text{End}_{\mathcal{Z}^s}(p'^* \mathcal{T}il_K))$. But this is impossible as we have seen in Lemma 2.13, there is an object with all Ext's with $p'^*(\mathbb{S}_\alpha \mathcal{Q}^*)$ vanishes but having non-trivial Ext with $p'^*(\mathbb{S}_\beta \mathcal{Q}^*)$.

More explicitly, we describe here the equivariant simple modules, considered as modules over $\text{End}_{\mathcal{O}^s_Z}(p^*(\oplus_\alpha \nabla_\alpha))$ and as representations.

Remark 5.9. As can be seen, in the set-up of this section, the conditions in Proposition 2.19 and Remark 2.20 are satisfied. Thus, $S_\beta = R\,\text{Hom}_{\mathcal{O}^s_Z}(p^*(\oplus_\alpha \nabla_\alpha), u_* \Delta_\beta)$ are all the equivariant simple objects over $\text{End}_{\mathcal{O}^s_Z}(p^*(\oplus_\alpha \nabla_\alpha))$. As has been remarked in 2.20, the finite dimensional algebra $\text{End}_{\mathcal{O}_{\text{Grass}}}(\oplus \nabla_\alpha) = \mathbb{C}QK(n, n-r)/I$ is a subring of $\text{End}_{\mathcal{O}^s_Z}(p^*(\oplus_\alpha \nabla_\alpha))$. The modules S_α are also simple modules over $\mathbb{C}QK(n, n-r)/I$. In fact, as $\mathbb{C}QK(n, n-r)/I$ is a finite dimensional basic algebra, S_α is the simple representation of the underlying quiver $QK(n, nn-r)$ corresponding to the vertex α. In this case, it is evident that S_α's are distinct as modules over $\text{End}_{\mathcal{O}^s_Z}(p^*(\oplus_\alpha \nabla_\alpha))$.

Let us look at the structure of S_β's as GL_n-representations. By definition,

$$\begin{aligned} S_\beta &= R\,\mathrm{Hom}_{\mathscr{O}_Z^s}(p^*(\oplus_\alpha \nabla_\alpha), u_*\Delta_\beta) \\ &\cong \mathrm{Hom}_{\mathscr{O}_Z^s}(p^*\nabla_\beta, u_*\Delta_\beta) \\ &\cong \mathrm{Hom}_{\mathrm{Grass}}(\nabla_\beta, \Delta_\beta) \\ &\cong H^{(n-r)r-|\alpha|}(\mathrm{Grass}, \mathbb{S}_\beta \mathcal{Q} \otimes \mathbb{S}_{(n-r)^r}\mathcal{Q}^* \otimes \mathbb{S}_{\alpha^{ct}}\mathcal{R}) \\ &\cong \mathbb{C}. \end{aligned}$$

All of them are 1-dimensional trivial representations.

The following Proposition is a direct consequence of Lemma 3.5.

Proposition 5.10. *Let S_α be as in Proposition 2.19. Then, the Ext's among them are given by*

$$\mathrm{Ext}^t(S_\alpha, S_\beta) \cong \oplus_s \oplus_{\lambda \in Q_{-1}(2s)} H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_{\beta^t}\mathcal{R}^* \otimes \mathbb{S}_{\alpha^t}\mathcal{R}),$$

where $Q_{-1}(2s) = \{\lambda \vdash 2s | \lambda = (a_1, \cdots, a_r | a_1 - 1, \cdots, a_r - 1)\}$ in the hook notation.

Note that the only element in $Q_{-1}(2)$ is $(2, 0, \cdots, 0)$ and the only element in $Q_{-1}(4)$ is $(3, 1, 0, \cdots, 0)$. One can easily calculate Ext^1 and Ext^2 with the aid of Lemma 3.9.

For $\lambda = (3, 1, 0, \cdots, 0)$, the $\gamma_{min(3,1,0,\cdots,0)}$ is given by Lemma 3.9 is (-2) if $n - r = 1$ with the corresponding $t_{min(3,1,0,\cdots,0)} = 2$; $\gamma_{min(3,1,0,\cdots,0)} = (-1, -2)$ if $n - r = 2$ with the corresponding $t_{min(3,1,0,\cdots,0)} = 2$; $\gamma_{min(3,1,0,\cdots,0)} = (-1, -1, -2)$ if $n - r \geq 3$ with the corresponding $t_{min(3,1,0,\cdots,0)} = 2$. Note that in any case above there is no operation described in Remark 3.10 satisfying the constrains given by Corollary 3.6.

For $\lambda = (2, 0, \cdots, 0)$, the $\gamma_{min(2,0,\cdots,0)}$ is given by Lemma 3.9 is (-1) if $n - r = 1$ with the corresponding $t_{min(2,0,\cdots,0)} = 1$; $\gamma_{min(2,0,\cdots,0)} = (0, \cdots, 0, -1, -1)$ if $n - r \geq 2$ with the corresponding $t_{min(2,0,\cdots,0)} = 1$.

Note that in the case $n - r = 1$ there is one operation described in Remark 3.10 satisfying the constrains given by Corollary 3.6, which gives $\gamma = (0)$ with $t = 2$. In the case $n - r = 2$ there is one operation described in Remark 3.10 satisfying the constrains given by Corollary 3.6 but can be applied successively, which gives $\gamma = (1, -1)$ and $\gamma = (2, -1)$ with the corresponding $t = 2$ and 3 respectively. In the case $n - r > 2$ there are a lot of operations. But if we only care about those with corresponding $t = 2$ and satisfying the constrains given by Corollary 3.6, there is only one which gives $\gamma = (1, 0, \cdots, 0, -1, -1)$.

The above analysis proves the following proposition. (Note that only $Q_{-1}(2s)$ with $s = 0, 1, 2$ contributes to Ext^1 and Ext^2.)

Proposition 5.11. *Let S_α be as in Proposition 2.19. We have*

$$\mathrm{Ext}^1(S_\alpha, S_\beta) \cong \begin{cases} (C^\alpha_{\beta,(1,0,\cdots,0)} E^*) \oplus (C^\beta_{\alpha,(1,0,\cdots,0)} E^*), & \text{if } n - r = 1 \\ (E^*) \oplus (C^\alpha_{\beta,(1,0,\cdots,0)} \mathbb{C}), & \text{if } n - r \geq 2 \end{cases}$$

$$\mathrm{Ext}^2(S_\alpha, S_\beta) \cong \begin{cases} (C^\alpha_{\beta,(1,1,0,\cdots,0)} \mathbb{S}_2 E^*) \oplus (C^\beta_{\alpha,(1,1,0,\cdots,0)} \mathbb{S}_2 E^*) \oplus (\delta^\beta_\alpha \wedge^2 E^*), \\ \text{if } n - r = 1 \\ (C^{\beta^t}_{\alpha^t,(1,-1)} \wedge^2 E^*) \oplus (C^{\beta^t}_{\alpha^t,(-1,-2)} E^*) \oplus (C^\beta_{\alpha,(1,1,0,\cdots,0)} \mathbb{S}_2 E^*) \oplus (C^\beta_{\alpha,(2,0,\cdots,0)} \wedge^2 E^*), \\ \text{if } n - r = 2 \\ (C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-1,-2)} \mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-1,-1)} E^*) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E^*) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E^*), \\ \text{if } n - 2 \geq 2 \end{cases}$$

5.3. **Maximal minors.** We assume $r = n - 1$, $S^s = \mathbb{C}[x_{i,j}]_{1\le i\le j\le n}$ and $R^s = S^s/(\det(x_{ij}))$. In this case, R^s is always Gorenstein, since it's a hypersurface.

The inverse image of the tilting bundle from the Grassmannian is still a tilting bundle by the same argument as before. More explicitly, it is

$$\bigoplus_{i=0}^{n-1} \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q).$$

Proposition 5.12. *The endomorphism ring* $\mathrm{End}_S(q'_* p'^* \mathscr{T}il_K)$ *is isomorphic to the path algebra of the quiver:*

$$\begin{tikzcd}
\bullet_0 \arrow[r, bend left=60, "\alpha_1"] \arrow[r, bend left=45, "\alpha_2"] \arrow[r, bend left=35, phantom, "\cdots"] \arrow[r, bend left=20, "\alpha_n"] & \bullet_1 \arrow[l, bend left=20, "\beta_n"] \arrow[l, bend left=35, phantom, "\cdots"] \arrow[l, bend left=45, "\beta_2"] \arrow[l, bend left=60, "\beta_1"] \arrow[r, bend left=60, "\alpha_1"] \arrow[r, bend left=45, "\alpha_2"] \arrow[r, bend left=35, phantom, "\cdots"] \arrow[r, bend left=20, "\alpha_n"] & \cdots \arrow[l, bend left=20, "\beta_n"] \arrow[l, bend left=35, phantom, "\cdots"] \arrow[l, bend left=45, "\beta_2"] \arrow[l, bend left=60, "\beta_1"] \arrow[r, bend left=60, "\alpha_1"] \arrow[r, bend left=45, "\alpha_2"] \arrow[r, bend left=35, phantom, "\cdots"] \arrow[r, bend left=20, "\alpha_n"] & \bullet_{n-1} \arrow[l, bend left=20, "\beta_n"] \arrow[l, bend left=35, phantom, "\cdots"] \arrow[l, bend left=45, "\beta_2"] \arrow[l, bend left=60, "\beta_1"]
\end{tikzcd}$$

with relations:

$$\begin{gathered}
\alpha_i\alpha_j + \alpha_j\alpha_i,\\
\beta_i\beta_j + \beta_j\beta_i,\\
(\alpha_i\beta_j + \beta_j\alpha_i) - (\alpha_j\beta_i + \beta_i\alpha_j),
\end{gathered}$$

where for any term not making sense at some vertex, it is to be understand as dropped.

In the language of equivariant quivers, the above quiver can be written as

$$\begin{tikzcd}
\bullet_0 \arrow[r, bend left, "\alpha_0(E)"] & \bullet_1 \arrow[l, bend left, "\beta_0(E)"] \arrow[r, bend left, "\alpha_1(E)"] & \cdots \arrow[l, bend left, "\beta_1(E)"] \arrow[r, bend left, "\alpha_{n-1}(E)"] & \bullet_{n-1} \arrow[l, bend left, "\beta_{n-2}(E)"]
\end{tikzcd}$$

and the relations:

$$\begin{gathered}
\alpha_i\alpha_{i+1}(\wedge^2 E);\\
\beta_{i+1}\beta_i(\wedge^2 E);\\
(\beta_i\alpha_i + \alpha_{i+1}\beta_{i+1})(\mathbb{S}_2 E).
\end{gathered}$$

Proof. See the proof of Proposition 5.14 and the proof of Proposition 5.15. ◻

A description of the Ext's between the simples in the module category over the path algebra of quiver with relations is given by Proposition 5.10.

Example 5.13. For any integer $a \in [0, n-1]$, the minimal projective resolution of the simple object S_a is:

$$\begin{tikzcd}
P_a & \begin{matrix} E\otimes P_{a-1} \\ \oplus \\ E\otimes P_{a+1} \end{matrix} \arrow[l] & \begin{matrix} \mathbb{S}_2E\otimes P_{a-2} \\ \oplus \\ \wedge^2E\otimes P_a \\ \oplus \\ \mathbb{S}_2E\otimes P_{a+2} \end{matrix} \arrow[l] & \begin{matrix} \mathbb{S}_3E\otimes P_{a-3} \\ \oplus \\ \mathbb{S}_{22}E\otimes P_{a-2} \\ \oplus \\ \mathbb{S}_{211}E\otimes P_a \\ \oplus \\ \mathbb{S}_{22}E\otimes P_{a+2} \\ \oplus \\ \mathbb{S}_3E\otimes P_{a+3} \end{matrix} \arrow[l] & \begin{matrix} \mathbb{S}_4E\otimes P_{a-4} \\ \oplus \\ \mathbb{S}_{32}E\otimes P_{a-3} \\ \oplus \\ \mathbb{S}_{221}E\otimes P_{a-1} \\ \oplus \\ \mathbb{S}_{221}E\otimes P_{a+1} \\ \oplus \\ \mathbb{S}_{32}E\otimes P_{a+3} \\ \oplus \\ \mathbb{S}_4E\otimes P_{a+4} \end{matrix} \arrow[l]
\end{tikzcd}.$$

If a is close to the boundary (i.e., 0 and n-1), some terms in the above resolution doesn't exist. In those cases, those terms should be understood as dropped.

5.4. **Higher codimension cases.** In higher codimension cases, the tilting bundle is given by $\oplus_{\lambda\subset(2,2)} p'^*\mathbb{S}_\lambda Q^*$. And then we show the path algebra is derived equivalent to the category of quasi-coherent sheaves over the commutative desingularization, i.e., the path algebra is a non-commutative desingularization. We also prove that the commutative desingularization gives the fine moduli space of certain bound representations of this quiver.

Now we start using Proposition 5.11 to describe the quiver with relations for this non-commutative desingularization.

Proposition 5.14. *Assume $n-r \geq 2$. In the quiver with relations for the algebra $\Lambda = \mathrm{End}_{\mathscr{O}_{\mathcal{Z}^s}}(p'^*\mathscr{T}il_K)$ as in Proposition 2.19, the vertices correspond to the sub-partitions of $(n-r)^r$, and the arrow from β to α is given by E if $C^{\alpha}_{\beta,(1,0,\cdots,0)} \neq 0$ and given by $\mathbb{C}$ if $C^{\beta}_{\alpha,(1,1,0,\cdots,0)} \neq 0$. No arrows otherwise.*

Proposition 5.15. *In the quiver with relations for the algebra $\Lambda = \mathrm{End}_{\mathscr{O}_{\mathcal{Z}^s}}(p'^*\mathscr{T}il_K)$ as in Proposition 2.19, the relations are generated by the following sub-representations of the arrows in $\mathrm{Hom}(\beta,\alpha)$:*

- $$(C^{\beta^t}_{\alpha^t,(1,-1)} \wedge^2 E) \oplus (C^{\beta^t}_{\alpha^t,(-1,-2)} E) \oplus (C^{\beta}_{\alpha,(1,1,0,\cdots,0)} \mathbb{S}_2 E) \oplus (C^{\beta}_{\alpha,(2,0,\cdots,0)} \wedge^2 E)$$
  in the case $n-r=2$;
- $$(C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-1,-2)} \mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-1,-1)} E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E)$$
  in the case $n-r \geq 3$.

Similar to Proposition 4.10, we have the following equivalence of equivariant derived categories.

Proposition 5.16. *The functor $R\,\mathrm{Hom}_{\mathscr{O}_{\mathcal{Z}^s}}(p'^*\mathscr{T}il_K, -)$ induces an equivalence between $D^b_G(Coh(\mathcal{Z}^s))$ and $D^b_G(\mathrm{End}_{\mathscr{O}_{\mathcal{Z}^s}}(p'^*\mathscr{T}il_K)\text{-}mod)$.*

5.5. **Examples.** We look at two examples with $r = n-1$, $S^s = \mathbb{C}[x_{i,j}]_{1\leq i\leq j\leq n}$ and $R^s = S^s/(det)$.

The inverse image of the tilting bundle is

$$\bigoplus_{i=0}^{n-1} \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q).$$

The presentations of the global sections of it's direct summands as S-modules are given by

$$0 \leftarrow H^0(\mathcal{Z}^s, \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q)) \leftarrow \mathbb{S}_{(1^i,0^{n-i})}E^* \otimes S \leftarrow \mathbb{S}_{(2^i,1^{n-i})}E^* \otimes S.$$

Let's take $n = 3$ and $r = 2$ in this section for a concrete example. Here $S^s = \mathbb{C}[x_{i,j}]_{1\leq i\leq j\leq 3}$ and $R^s = S^s/(det)$.

In this example, R is a normal Gorenstein domain since it is a hypersurface ring. Consequently, if $\mathrm{End}_{\mathcal{Z}^s}(p'^*\mathscr{T}il_K)$ is maximal Cohen-Macaulay and $q'_*p'^*\mathscr{T}il_K$ is reflexive, then $\mathrm{End}_{\mathcal{Z}^s}(p'^*\mathscr{T}il_K)$ is a non-commutative crepant desingularization due to the following proposition.

The inverse image of the tilting bundle consists of three direct summands: $\mathrm{Sym}(\mathbb{S}_2 Q)$, $Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q)$, $\wedge^2 Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q)$. Let's denote their global sections as R-modules, (and S^s-modules automatically), by $M_i := H^0(\mathcal{Z}^s, \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q))$ with $i = 0,1,2$. The presentations of their global sections as S^s-modules are given by

$$0 \leftarrow H^0(\mathcal{Z}^s, \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q)) \leftarrow \mathbb{S}_{(1^i,0^{3-i})}E^* \otimes S^s \leftarrow \mathbb{S}_{(2^i,1^{3-i})}E^* \otimes S^s.$$

The Hom's between them are

$$\mathrm{Hom}_R(M_i, M_j) = H^0(\mathcal{Z}^s, \wedge^j Q^* \otimes \wedge^i Q \otimes \mathrm{Sym}(\mathbb{S}_2 Q)).$$

More explicitly,

- $\mathrm{Hom}_R(M_0, M_i) = M_i$,
- $\mathrm{Hom}_R(M_i, M_0) = M_i$,
- $\mathrm{Hom}_R(M_2, M_i) = M_{2-i}$,
- $\mathrm{Hom}_R(M_i, M_2) = M_{2-i}$.

Through some computations, we get

$$\mathrm{Hom}_R(M_1, M_1) = M_0 \oplus N$$

and the presentation of N is

$$0 \leftarrow N \leftarrow \mathbb{S}_{(200)}E^* \oplus \mathbb{S}_{(110)}E^* \otimes S^s \leftarrow \mathbb{S}_{(220)}E^* \oplus \mathbb{S}_{(211)}E^* \otimes S^s.$$

The endomorphism ring is isomorphic to the path algebra of the following quiver with relations.

The quiver is



with relations:

$$\gamma_i\alpha_j + \gamma_j\alpha_i,$$

$$\beta_j\delta_i + \beta_i\delta_j,$$

$$\beta_j\alpha_i - \beta_i\alpha_j,$$

$$\gamma_j\delta_i - \gamma_i\delta_j,$$

$$(\alpha_i\beta_j + \delta_j\gamma_i) - (\alpha_j\beta_i + \delta_i\gamma_j).$$

The Hom between any two direct summands is graded, with grading given by the weight of $\mathbb{G}_m$-action on E. The grading defined this way is different but (strictly) finer than the one used in the proof of Proposition 2.19.

The Hilbert series of those modules can be computed from the presentations:

- $H_{M_0} = \frac{1-t^6}{(1-t^2)^6}$;
- $H_{M_1} = \frac{3-3t^4}{(1-t^2)^6}$;
- $H_{M_2} = \frac{3-3t^2}{(1-t^2)^6}$;
- $H_N = \frac{9-9t^2}{(1-t^2)^2}$.

Putting the Hilbert series of the Hom's into a matrix, we get

$$\frac{1}{(1-t^2)^5} \times \begin{pmatrix} 1+t^2+t^4 & 3t+3t^3 & 3t^2 \\ 3t+3t^3 & 1+10t^2+t^4 & 3t+3t^3 \\ 3t^2 & 3t+3t^3 & 1+t^2+t^4 \end{pmatrix}$$

The coefficients in front of each monomials in the entries of the inverse matrix gives the multiplicity of the projectives in the resolution of the simples. The matrix above has an inverse with polynomial entries, which corresponds to the fact that the derived category over the endomorphism ring has finite global dimension.

The inverse matrix is

$$\begin{pmatrix} -t^6-3t^4+3t^2+1 & 3t^5-3t & -6t^4+6t^2 \\ 3t^5-3t & -t^6-3t^4+3t^2+1 & 3t^5-3t \\ -6t^4+6t^2 & 3t^5-3t & -t^6-3t^4+3t^2+1 \end{pmatrix}$$

It's easy to guess the resolution of the simples from this matrix. It is easy to verify that this resolution coincide with the one given by Lemma 2.26 and Proposition 5.10.

$$S_0: \quad P_0 \leftarrow E\otimes P_1 \leftarrow \begin{array}{c} \wedge^2 E\otimes P_0 \\ \oplus \\ \mathbb{S}_2 E\otimes P_2 \end{array} \leftarrow \begin{array}{c} \mathbb{S}_{211} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{22} E\otimes P_2 \end{array} \leftarrow \mathbb{S}_{221} E\otimes P_1 \leftarrow \mathbb{S}_{222} E\otimes P_0 \leftarrow 0\ ;$$

$$S_1: \quad P_1 \leftarrow \begin{array}{c} E\otimes P_0 \\ \oplus \\ E\otimes P_2 \end{array} \leftarrow \wedge^2 E\otimes P_1 \leftarrow \mathbb{S}_{211} E\otimes P_1 \leftarrow \begin{array}{c} \mathbb{S}_{221} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{221} E\otimes P_2 \end{array} \leftarrow \mathbb{S}_{222} E\otimes P_1 \leftarrow 0\ .$$

Example 5.17. Let's look at one more example with $n = 4$ and $r = 3$.

As before, the direct summands of the tilting bundle over the desingularization are $\wedge^i Q^* \otimes \mathrm{Sym}_2(\mathbb{S}_2 Q)$ where $i = 0, \dots, 3$.

The presentations of their global sections as S^s-modules are given by

$$0 \leftarrow M_i := H^0(\mathcal{Z}, \wedge^i Q^* \otimes \mathrm{Sym}(\mathbb{S}_2 Q)) \leftarrow \mathbb{S}_{(1^i,0^{n-i})} E^* \otimes S^s \leftarrow \mathbb{S}_{(2^i,1^{n-i})} E^* \otimes S^s.$$

The endomorphism ring is isomorphic to the path algebra of the quiver:



with relations:

$$\alpha_i\alpha_j + \alpha_j\alpha_i,$$
$$\beta_i\beta_j + \beta_j\beta_i,$$
$$(\alpha_i\beta_j + \beta_j\alpha_i) - (\alpha_j\beta_i + \beta_i\alpha_j).$$

To get the Hilbert polynomials of the Hom's among them, we only need to compute the presentations of two modules, i.e., $\mathrm{Hom}(M_1, M_1)$ and $\mathrm{Hom}(M_1, M_2)$. They are given as follows.

- $\mathrm{Hom}(M_1, M_1) = M_0 \oplus C_1$, where
  $$0 \leftarrow C_1 \leftarrow \mathbb{S}_{11} E\otimes S^s \oplus \mathbb{S}_2 E\otimes S^s \leftarrow \mathbb{S}_{2211} E\otimes S^s \oplus \mathbb{S}_{222} E\otimes S^s$$
  is exact;
- $\mathrm{Hom}(M_1, M_2) = M_1 \oplus C_2$, where
  $$0 \leftarrow C_2 \leftarrow \mathbb{S}_{111} E\otimes S^s \oplus \mathbb{S}_{21} E\otimes S^s \leftarrow \mathbb{S}_{221} E\otimes S^s \oplus \mathbb{S}_{2111} E\otimes S^s$$
  is exact.

The matrix of Hilbert polynomials between their Hom's is

$$\frac{1}{(1-t^2)^{10}} \begin{pmatrix} 1-t^8 & 4-4t^6 & 6-6t^4 & 4-4t^2 \\ 4-4t^6 & 17-16t^4-t^8 & 28-24t^2-4t^6 & 6-6t^4 \\ 6-6t^4 & 28-24t^2-4t^6 & 17-16t^4-t^8 & 4-4t^6 \\ 4-4t^2 & 6-6t^4 & 4-4t^6 & 1-t^8 \end{pmatrix}.$$

It's inverse matrix is

$$\begin{pmatrix} 15t^8-6t^{10}-t^{12}-15t^4+6t^2+1 & 20t^5-20t^7+4t^{11}-4t & -20t^4+20t^8-10t^{10}+10t^2 & -20t^3+60t^5-60t^7+20t^9 \\ 20t^5-20t^7+4t^{11}-4t & 15t^8-6t^{10}-t^{12}-15t^4+6t^2+1 & 20t^5-20t^7+4t^{11}-4t & -20t^4+20t^8-10t^{10}+10t^2 \\ -20t^4+20t^8-10t^{10}+10t^2 & 20t^5-20t^7+4t^{11}-4t & 15t^8-6t^{10}-t^{12}-15t^4+6t^2+1 & 20t^5-20t^7+4t^{11}-4t \\ -20t^3+60t^5-60t^7+20t^9 & -20t^4+20t^8-10t^{10}+10t^2 & 20t^5-20t^7+4t^{11}-4t & 15t^8-6t^{10}-t^{12}-15t^4+6t^2+1 \end{pmatrix}.$$

One can guess the resolution of the simples from this matrix. Again, it is easy to verify that this resolution coincide with the one given by Lemma 2.26 and Proposition 5.10.

$$S_0: \quad P_0 \longleftarrow E\otimes P_1 \longleftarrow \begin{matrix} \wedge^2 E\otimes P_0 \\ \oplus \\ \mathbb{S}_2 E\otimes P_2 \end{matrix} \longleftarrow \begin{matrix} \mathbb{S}_3 E\otimes P_3 \\ \oplus \\ \mathbb{S}_{211} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{22} E\otimes P_2 \end{matrix} \longleftarrow \begin{matrix} \mathbb{S}_{221} E\otimes P_1 \\ \oplus \\ \mathbb{S}_{32} E\otimes P_3 \end{matrix}$$

$$\longleftarrow \begin{matrix} \mathbb{S}_{3211} E\otimes P_1 \\ \oplus \\ \mathbb{S}_{331} E\otimes P_3 \end{matrix} \leftarrow \begin{matrix} \mathbb{S}_{333} E\otimes P_3 \\ \oplus \\ \mathbb{S}_{3221} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{3311} E\otimes P_2 \end{matrix} \leftarrow \begin{matrix} \mathbb{S}_{3322} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{3331} E\otimes P_2 \end{matrix} \leftarrow \mathbb{S}_{3332} E\otimes P_1 \leftarrow \mathbb{S}_{3333} E\otimes P_0 \leftarrow 0;$$

$$S_1: \quad P_1 \longleftarrow \begin{matrix} E\otimes P_0 \\ \oplus \\ E\otimes P_2 \end{matrix} \longleftarrow \begin{matrix} \wedge^2 E\otimes P_1 \\ \oplus \\ \mathbb{S}_2 E\otimes P_3 \end{matrix} \longleftarrow \begin{matrix} \mathbb{S}_{211} E\otimes P_1 \\ \oplus \\ \mathbb{S}_{22} E\otimes P_3 \end{matrix} \longleftarrow \begin{matrix} \mathbb{S}_{221} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{221} E\otimes P_2 \end{matrix}$$

$$\longleftarrow \begin{matrix} \mathbb{S}_{3211} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{3211} E\otimes P_2 \end{matrix} \leftarrow \begin{matrix} \mathbb{S}_{3221} E\otimes P_1 \\ \oplus \\ \mathbb{S}_{3311} E\otimes P_3 \end{matrix} \leftarrow \begin{matrix} \mathbb{S}_{3322} E\otimes P_1 \\ \oplus \\ \mathbb{S}_{3331} E\otimes P_3 \end{matrix} \leftarrow \begin{matrix} \mathbb{S}_{3332} E\otimes P_0 \\ \oplus \\ \mathbb{S}_{3332} E\otimes P_2 \end{matrix} \leftarrow \mathbb{S}_{3333} E\otimes P_1 \leftarrow 0.$$

Now let's look at an example of higher codimension symmetric minors case. Let's take $n = \dim E = 4$ and $r = 2$.

The first two steps are relatively easy. By direct computation, we get the following quiver (recall that vertices are indexed by Young diagrams):



with relations:

- Hom($\ydiagram{2}$, ∅): $\alpha_1\alpha_3(\wedge^2 E)$;
- Hom($\ydiagram{1,1}$, ∅): $\alpha_1\alpha_2(\mathbb{S}_2 E)$;
- Hom($\ydiagram{2,2}$, $\ydiagram{2}$): $\alpha_5\alpha_6(\wedge^2 E)$;
- Hom($\ydiagram{2,2}$, $\ydiagram{1,1}$): $\alpha_4\alpha_6(\mathbb{S}_2 E)$.
- Hom($\ydiagram{2,1}$, $\ydiagram{1}$): $\alpha_2\alpha_4 - \alpha_3\alpha_5(\mathbb{S}_2 E \oplus \wedge^2 E)$;
- Hom($\ydiagram{1}$, $\ydiagram{1}$): $\alpha_3\beta_1\alpha_1 - \alpha_3\alpha_5\beta_2(\wedge^2 E)$;

- Hom($\ydiagram{2,1}$, $\ydiagram{2,1}$): $\alpha_6\beta_3\alpha_5 - \beta_2\alpha_3\alpha_5(\wedge^2 E)$;
- Hom($\varnothing$, $\ydiagram{2,1}$): $\beta_2\alpha_3\beta_1 - \alpha_6\beta_3\beta_1(E)$;
- Hom($\ydiagram{1}$, $\ydiagram{2,2}$): $\beta_3\beta_1\alpha_1 - \beta_3\alpha_5\beta_2(E)$.

Using Olver's description of Pieri inclusions, (see, e.g., 1.2 of [SW10],) it's easy to check the above listed subrepresentations acts trivially on $p'^* \oplus \mathbb{S}_\alpha \mathcal{Q}^*$. Then Proposition 5.15 yields that these are all the relations.

6. Pfaffian varieties of anti-symmetric matrices

What we will do in this section is parallel to the previous section.

6.1. **Review of the commutative desingularization.** Let E be a vector spaces over k of dimension n and H^a be the subspace of $\operatorname{Hom}_k(E^*, E)$ consisting of anti-symmetric morphisms. This space can be identified with $\wedge^2(E)$. Upon choosing a set of basis, H^s can be identified with the set of symmetric $(n\times n)$-matrices (x_{ij}) with $x_{ij} = -x_{ji}$, whose coordinate ring can be identified with $S^a = k[x_{ij}]_{i<j}$.

We get a universal morphism $\varphi : \mathcal{E}^* \to \mathcal{E}$ over H^a. For $r \leq n$, we desingulariz the locus $\operatorname{Spec} R^a$, where φ has rank $\leq r$. In other words, R^a is the quotient of S^a by the ideal generated by the $(r+1)\times(r+1)$ minors of (x_{ij}).

Let

$$0 \to \mathcal{R} \to E \times \operatorname{Grass} \to \mathcal{Q} \to 0$$

be the tautological sequence over Grass, where Grass is the grassmannian of $(n-r)$-planes in E.

We will add an upper script a for the varieties and keep the same notations for the maps. The incidence variety $\mathcal{Z}^a$ of $\mathcal{Y}^a$ desingularizing $\operatorname{Spec} R^a$, is defined by $\mathcal{Z}^a = \{(g,h) \in \operatorname{Grass} \times H^a : \operatorname{im} h \in g\}$. We use the following diagram to illustrate all the notations.

$$\begin{tikzcd}
\mathcal{Z}^a \arrow[dr, hook, "j"] \arrow[drr, bend left, "p'"] \arrow[dd, "q'"'] & & \\
& \mathcal{Y}^a = \operatorname{Grass} \times H^a \arrow[r, "p"] \arrow[d, "q"'] & \operatorname{Grass} = \operatorname{Grass}_{n-r}(E) \\
\operatorname{Spec} R^a \arrow[r, hook, "i"] & H^a = \wedge^2(E^*) &
\end{tikzcd}$$

Proposition 6.1 (6.4.2 in [W03])**.** *The variety $\mathcal{Z}^a$ is a desingularization of* $\operatorname{Spec} R^a$.

The variety $\mathcal{Z}^a$ can be described as the total space of the vector bundle $\wedge^2(\mathcal{Q}^*)$ over Grass. Equivalently, $p' : \mathcal{Z}^a \to \operatorname{Grass}$ is an affine morphism with $p_*\mathcal{O}_{\mathcal{Z}^a}$ equal to the sheaf of algebra $\operatorname{Sym}(\wedge^2 \mathcal{Q})$.

6.2. **A tilting bundle over the desingularization.** As before, we pull back $\mathcal{T}il_K = \oplus_{\lambda\subset(n-r)^r} \mathbb{S}_\lambda \mathcal{Q}^*$, the Kapranov's tilting bundle over $\operatorname{Grass}_{n-r}(E)$, by $p' : \mathcal{Z}^a \to \operatorname{Grass}$. Similar to the symmetric case, we can show that for all $i > 0$, and all α, β sub-partitions of $((n-r)^r)$,

$$H^i(\mathcal{Z}^a, p'^* \mathcal{H}om_{\mathcal{O}_{\operatorname{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*)) = 0.$$

From the claim above, using Lemma2.6 and Lemma2.5, we get the following

Proposition 6.2. *The Kapranov's tilting bundle over* $\mathrm{Grass}_{n-r}(F)$ *is denoted by* $\mathcal{T}il_K$*. The rank* r *determinantal variety* $\operatorname{Spec} R^a$ *of anti-symmetric matrices and its desingularization* $\mathcal{Z}^a$ *are as above. The bundle* $p'^*\mathcal{T}il_K$ *is a tilting bundle over* $\mathcal{Z}^a$*.*

Lemma 6.3. *The quasi-coherent sheaf* $\mathrm{End}_{\mathcal{Z}^a}(p'^*\mathcal{T}il_K)$ *over* $\operatorname{Spec} R^a$ *is never maximal Cohen-Macaulay.*

Proof. According to Lemma 2.9, it suffices to compute $H^i(\mathrm{Grass}, \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^a} \otimes \mathrm{Sym}(\wedge^2 \mathcal{Q}))$ for $i > 1$ and α, $\beta \subset (n-r)^r$. The sheaf $\omega_{\mathcal{Z}^a}$ has been computed in 6.7 of [W03], which says $\omega_{\mathcal{Z}^a} \cong \mathbb{S}_{(-n-1+r)^n} E^* \otimes \mathbb{S}_{(n-r-1)^r} \mathcal{Q}^*$. Hence,

$$\mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^a} \otimes \mathrm{Sym}(\wedge^2 \mathcal{Q}) \cong \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \mathbb{S}_{(-n-1+r)^n} E^* \otimes \mathbb{S}_{(n-r-1)^r} \mathcal{Q}^* \otimes \mathrm{Sym}(\wedge^2 \mathcal{Q}).$$

Using Cauchy-Littlewood and Littlewood-Richardson formulas, as well as Bott's Theorem, we see that

$$H^i(\mathrm{Grass}, \mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \otimes \omega_{\mathcal{Z}^a} \otimes \mathrm{Sym}(\wedge^2 \mathcal{Q})) = 0$$

if and only if $\mathbb{S}_\alpha \mathcal{Q}^* \otimes \mathbb{S}_\beta \mathcal{Q} \cong \oplus \mathbb{S}_\lambda \mathcal{Q}^*$ has $\lambda_1 < 1$ for all λ. This can never happen for all α, $\beta \subset (n-r)^r$ for any n or r. □

By Theorem 7.6 in [HV07], since $\mathcal{Z}^a$ is smooth, $\mathrm{End}_{\mathcal{Z}^a}(p'^*\mathcal{T}il_K)$ has finite global dimension. Using Proposition 2.7, we get the following.

Proposition 6.4. *If* $r > 2$*, the map*

$$\mathrm{End}_{\mathcal{Z}^a}(p'^*\mathcal{T}il_K) \to \mathrm{End}_S(q'_* p'^* \mathcal{T}il_K)$$

is an isomorphism of rings. In particular, $\mathrm{End}_{\mathcal{Z}^a}(p'^*\mathcal{T}il_K)$ *is a non-commutative desingularization (but never crepant).*

Now, we describe the Ext's between the equivariant simples. Then, we will use that to get the shape of quivers with relations of the non-commutative desingularization. As similar things has been done in the symmetric case, we will omit some details in the proofs.

Proposition 6.5. *Let* S_α *be as in Proposition 2.19. Then, the Ext's among them are given by*

$$\mathrm{Ext}^t(S_\alpha, S_\beta) \cong \oplus_s \oplus_{\lambda \in Q_1(2s)} H^{t-s-|\beta|+|\alpha|}(\mathrm{Grass}, \mathbb{S}_\lambda \mathcal{Q}^* \otimes \mathbb{S}_{\beta^t} \mathcal{R}^* \otimes \mathbb{S}_{\alpha^t} \mathcal{R}),$$

where $Q_1(2s) = \{\lambda \vdash 2s \mid \lambda = (a_1, \cdots, a_r | a_1 + 1, \cdots, a_r + 1)\}$ *in the hook notation.*

Note that the only element in $Q_1(2)$ is $(1, 1, 0, \cdots, 0)$ and the only element in $Q_1(4)$ is $(2, 1, 1, 0, \cdots, 0)$. One can easily calculate Ext^1 and Ext^2 with the aid of Lemma 3.9.

For $\lambda = (1, 1, 0, \cdots, 0)$, the $\gamma_{min(1,1,0,\cdots,0)}$ is given by Lemma 3.9 is $(0, \cdots, 0, -2)$ with the corresponding $t_{min(1,1,0,\cdots,0)} = 1$. There are a lot of operations described in Remark 3.10, but if we only care about those with corresponding $t = 2$ and satisfying the constrains given by Corollary 3.6, there is only one which gives $\gamma = (1, 0, \cdots, 0, -2)$.

For $\lambda = (2, 1, 1, 0, \cdots, 0)$, the $\gamma_{min(2,1,1,0,\cdots,0)}$ is given by Lemma 3.9 is (-3) if $n - r = 1$ with the corresponding $t_{min(2,1,1,0,\cdots,0)} = 2$; $\gamma_{min(2,1,1,0,\cdots,0)} = (0, \cdots, 0, -1, -3)$ if $n - r \geq 2$ with the corresponding $t_{min(2,1,1,0,\cdots,0)} = 2$.

The above analysis proves the following proposition. (Note that only $Q_1(2s)$ with $s = 0, 1, 2$ contributes to Ext^1 and Ext^2.)

Proposition 6.6. *Let* S_α *be as in Proposition 2.19. We have*

$$\mathrm{Ext}^1(S_\alpha, S_\beta) \cong (C^\alpha_{\beta,(1,0,\cdots,0)} E^*) \oplus (C^\beta_{\alpha,(1,1,0,\cdots,0)} \mathbb{C});$$

$$\mathrm{Ext}^2(S_\alpha, S_\beta) \cong \begin{cases} (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0)} \wedge^3 E^*) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E^*), \\ \mathit{if}\; n-r=1 \\ (C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-3)} \mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-2)} E^*) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E^*) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E^*), \\ \mathit{if}\; n-r \geq 2 \end{cases}$$

Now we start using Proposition 6.6 to describe the quiver with relations for this non-commutative desingularization.

Proposition 6.7. *In the quiver with relations for the algebra* $\Lambda = \mathrm{End}_{\mathscr{O}_{\mathcal{Z}^a}}(p'^*\mathscr{T}il_K)$ *as in Proposition 2.19, the vertices correspond to the sub-partitions of* $(n-r)^r$*, and the arrow from* β *to* α *is given by* E *if* $C^{\alpha}_{\beta,(1,0,\cdots,0)} \neq 0$ *and given by* $\mathbb{C}$ *if* $C^{\beta}_{\alpha,(1,1,0,\cdots,0)} \neq 0$*. No arrows otherwise.*

Proposition 6.8. *In the quiver with relations for the algebra* $\Lambda = \mathrm{End}_{\mathscr{O}_{\mathcal{Z}^a}}(p'^*\mathscr{T}il_K)$ *as in Proposition 2.19, the relations are generated by the following sub-representations of the arrows in* $\mathrm{Hom}(\beta, \alpha)$*:*

- $$(C^{\beta^t}_{\alpha^t,(1,0,\cdots,0)} \wedge^3 E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E)$$
  in the case $n-r=1$*;*
- $$(C^{\beta^t}_{\alpha^t,(0,\cdots,0,-1,-3)} \mathbb{C}) \oplus (C^{\beta^t}_{\alpha^t,(1,0,\cdots,0,-2)} E) \oplus (C^{\beta^t}_{\alpha^t,(2,0,\cdots,0)} \mathbb{S}_2 E) \oplus (C^{\beta^t}_{\alpha^t,(1,1,0,\cdots,0)} \wedge^2 E)$$
  in the case $n-r \geq 2$*.*

Again, we have the following equivalence of equivariant derived categories.

Proposition 6.9. *The functor* $R\,\mathrm{Hom}_{\mathscr{O}_{\mathcal{Z}^a}}(p'^*\mathscr{T}il_K, -)$ *induces an equivalence*

$$D^b_G(Coh(\mathcal{Z}^a)) \cong D^b_G(\mathrm{End}_{\mathscr{O}_{\mathcal{Z}^a}}(p'^*\mathscr{T}il_K)\text{-}mod).$$

7. DETERMINANTAL VARIETIES

In this section we study a non-commutative desingularization of determinantal varieties in the space of matrices.

7.1. **Review of the commutative desingularization.**

Now we start describing a desingularization of determinantal varieties. We will follow the notations in the paper by Buchweitz, et al [BLV09].

Let G and$F$ be vector spaces of dimension m, n respectively with $m \geq n$, and $H = \mathrm{Hom}_k(G, F)$. Upon choosing a set of basis $f_1, \cdots, f_m$ and $g_1, \cdots, g_n$, H is identified with the set of $(m \times n)$-matrices (x_{ij}). The coordinate ring of H can be identified with $S = k[x_{ij}]$.

By base extension from $\mathrm{Spec}\, k$ to H, we get two vector bundles $\mathcal{F}$ and $\mathcal{G}$ over H, and a universal morphism $\varphi : \mathcal{G} \to \mathcal{F}$. For $r \leq n$, we desingularize the locus $\mathrm{Spec}\, R$, where φ has rank $\leq r$. In other words, R is the quotient of S by the ideal generated by the $(r+1)\times(r+1)$ minors of (x_{ij}).

We take the grassmannian of $n-r$ planes in F^*, denoted by Grass. Let $\mathcal{Y}$ be the product Grass $\times H$, p and q be the projection to Grass and H respectively. Inside of $\mathcal{Y}$, there is an incidence variety, denoted by $\mathcal{Z}$, defined by

$$\mathcal{Z} = \{(g, h) \in \mathrm{Grass} \times H : \mathrm{im}\, g \circ h = 0\}.$$

The inclusion $\mathcal{Z} \hookrightarrow \mathcal{Y}$ is denoted by j, and $\mathrm{Spec}\, R \to H$ by i. The induced map $\mathcal{Z} \to \mathrm{Spec}\, R$ from $q : \mathcal{Y} \to H$ is denoted by q', and $\mathcal{Z} \to \mathrm{Grass}$ from $p : \mathcal{Y} \to \mathrm{Grass}$ by p'.

These notations are summarized by the following diagram.

$$\begin{tikzcd}
\mathcal{Z} \arrow[dr, hook, "j"] \arrow[drr, bend left, "p'"] \arrow[dd, "q'"'] & & \\
& \mathcal{Y} = \mathrm{Grass} \times H \arrow[r, "p"] \arrow[d, "q"'] & \mathrm{Grass} = \mathrm{Grass}_{n-r}(F^*) \\
\operatorname{Spec} R \arrow[r, hook, "i"] & H = \mathrm{Hom}(G, F) &
\end{tikzcd}$$

Proposition 7.1 (6.1.1 in [W03])**.** *The variety* $\mathcal{Z}$ *is a desingularization of* $\operatorname{Spec} R$*.*

The variety $\mathcal{Z}$ can be described as the total space of a vector bundle over Grass.
Let

$$0 \to \mathcal{R} \to F^* \times \mathrm{Grass} \to \mathcal{Q} \to 0$$

be the tautological sequence over Grass. We apply the functor $G^* \otimes -$ to the dualized tautological sequence $0 \to \mathcal{Q}^* \to F \times \mathrm{Grass} \to \mathcal{R}^* \to 0$ to get

$$0 \to S \to E \to T \to 0$$

. Following [W03], let η be the sheaf of sections of $S^* = G \otimes \mathcal{Q}$. The desingularization $\mathcal{Z}$ is the total space of S. Equivalently, $p' : \mathcal{Z} \to \mathrm{Grass}$ is an affine morphism with $p_* \mathscr{O}_{\mathcal{Z}}$ equal to the sheaf of algebra $\mathrm{Sym}(\eta)$.

7.2. **A tilting bundle over the desingularization.** We will show, in this subsection, that $p'^* \mathscr{T}il_K$ is a tilting bundle over the desingularization $\mathcal{Z}$.

Now we prove the following

Proposition 7.2. *The Kapranov's tilting bundle over* $\mathrm{Grass}_{n-r}(F^*)$ *is denoted by*

$$\mathscr{T}il_K = \oplus_{\alpha \subset ((n-r)^r)} \mathbb{S}_\alpha \mathcal{Q}^*.$$

The rank r *determinantal variety* $\operatorname{Spec} R$ *and its desingularization* $\mathcal{Z}$ *are as above. The bundle* $p'^* \mathscr{T}il_K$ *is a tilting bundle over* $\mathcal{Z}$*.*

Proof. According to Theorem 2.7, it suffices to show that $H^i(\mathcal{Z}, p'^* \mathscr{H}om_{\mathscr{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*)) = 0$ for all $i > 0$, and all α, β sub-partitions of $((n-r)^r)$.

Use the Cauchy-Littlewood and the Littlewod-Richardson formulas, we can reduce the sheaf $p'^* \mathscr{H}om_{\mathscr{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*) \cong \mathscr{H}om_{\mathscr{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^* \otimes_{\mathscr{O}_{\mathrm{Grass}}} \mathrm{Sym}(\eta))$ into the form

$$\oplus (\mathbb{S}_\gamma \mathcal{Q}^*)^{\oplus C_\gamma} \otimes \mathbb{S}_\mu G$$

for some coefficients C_γ. By the Bott's theorem 3.2, all the higher cohomology of

$$p'^* \mathscr{H}om_{\mathscr{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^*) \cong \mathscr{H}om_{\mathscr{O}_{\mathrm{Grass}}}(\mathbb{S}_\alpha \mathcal{Q}^*, \mathbb{S}_\beta \mathcal{Q}^* \otimes_{\mathscr{O}_{\mathrm{Grass}}} \mathrm{Sym}(\eta))$$

vanishes. □

Proposition 7.3. *The module* $\mathrm{End}_{\mathcal{Z}}(p'^* \mathscr{T}il_K)$ *is reflexive and maximal Cohen-Macaulay.*

Proof. This has been proved in [BLV11]. □

Proposition 7.4. *The map*

$$\mathrm{End}_{\mathcal{Z}}(p'^* \mathscr{T}il_K) \to \mathrm{End}_S(q'_* p'^* \mathscr{T}il_K)$$

is an isomorphism of rings and it is a non-commutative desingularization of $\operatorname{Spec} R$*.*

Proof. By Proposition 2.7, it suffices to show the exceptional locus of q' has codimension at least 2 in both $\mathcal{Z}$ and $\operatorname{Spec} R$. The codimension of $\operatorname{Spec} R$ in $\operatorname{Spec} S$ is $\binom{n-r+1}{2}$, and hence the codimension of the singular locus of $\operatorname{Spec} R$ has codimension at least 2 unless $n = r$ in which case $\operatorname{Spec} R$ is smooth. The codimension of the exceptional locus in $\mathcal{Z}$ is the rank of the bundle $\mathcal{Z} \to \mathrm{Grass}$, which is at least 2 unless $m = \dim G = 1$ and $r = 1$ in which case $\operatorname{Spec} R$ is still smooth. □

By theorem 7.6 in [HV07], since $\mathcal{Z}$ is smooth, $\operatorname{End}_{\mathcal{Z}}(p'^{*}\mathcal{T}il_K)$ has finite global dimension.

7.3. **Formulas for the non-commutative desingularization.** Now we come to a formula for the noncommutative desingularization $\operatorname{End}_S(q'_{*}p'^{*}\mathcal{T}il_K) = \operatorname{End}_{\mathcal{Z}}(p'^{*}\mathcal{T}il_K)$.

Using a procedure similar to the one in Section 5, one can get the quiver with relations for the non-commutative desingularization. As the quiver with relations in this case is known to experts and yet to be published, instead, we explore the endomorphism ring using the basic theorem of geometric technique.

Let $\eta = \mathcal{Q} \otimes G$. The modules $H^0(\mathrm{Grass}, \mathrm{Sym}(\eta) \otimes \mathbb{S}_\alpha \mathcal{Q}^*)$ will be denoted by M_α.

In the basic theorem of geometric method 3.3, we take the vector bundle $\mathcal{V}$ to be $\mathbb{S}_\alpha\mathcal{V} = \mathbb{S}_\alpha\mathcal{Q}^*$, $V = \mathrm{Grass}$, and X to be H. By the same argument as in proposition 7.2, we get the vanishing of higher derived images for $M(\mathbb{S}_\alpha\mathcal{V})$, i.e.,

$$R^i q_* M(\mathbb{S}_\alpha\mathcal{V}) = R^i q_*(\mathcal{O}_{\mathcal{Z}} \otimes p^*\mathbb{S}_\alpha\mathcal{V}) = H^i(\mathrm{Grass}, \mathrm{Sym}(\eta) \otimes \mathbb{S}_\alpha\mathcal{V}) = 0$$

through the expansion of $\mathrm{Sym}(\eta) \otimes \mathbb{S}_\alpha\mathcal{V}$ using the Cauchy-Littlewood and the Littlewod-Richardson formulas, and Bott's theorem.

According to theorem 3.3, there is a presentation of M_α given by $F_{\alpha,1} \to F_{\alpha,0}$ with $F_{\alpha,i}$ defined to be

$$\begin{aligned}
F_{\alpha,i} &= \bigoplus_{j\geq 0} H^j(\mathrm{Grass}, \wedge^{i+j}(\xi) \otimes \mathbb{S}_\alpha\mathcal{Q}^*) \otimes S \\
&= \bigoplus_{j\geq 0} H^j(\mathrm{Grass}, \oplus_{|\mu|=i+j}\mathbb{S}_{\mu'}G \otimes \mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*) \otimes S \\
&= \bigoplus_{j\geq 0} \oplus_{|\mu|=i+j} H^j(\mathrm{Grass}, \mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*) \otimes \mathbb{S}_{\mu'}G \otimes S.
\end{aligned}$$

To compute the $F_{\alpha,i}$'s, all we need to do is to compute $H^j(\mathrm{Grass}, \mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*)$.

Theorem 7.5. *Notations as above,*

(1) $F_{\alpha,1} = \mathbb{S}_{(\alpha,1^{r+1-t},0^{m-r-1})}G \otimes \mathbb{S}_{(1^{r+1-t})}F^* \otimes S$;
(2) $F_{\alpha,0} = \mathbb{S}_\alpha G \otimes S$.

Proof. Assume $l(\alpha) = t$, i.e., $\alpha_t \neq 0$ and $\alpha_{t+1} = 0$, $|\mu| = j + 1$. To show (1), it suffices to prove that

(2)
$$H^j(\mathbb{S}_{\mu'}E \otimes \mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*) = \begin{cases} \mathbb{S}_{(\alpha,1^{r+1-t},0^{m-r-1})}G \otimes \mathbb{S}_{(1^{r+1-t})}F^*, \text{ if } j = |\alpha| + r - t - 1 \text{ and } \mu = (\alpha' + (r-t, 0^{n-r-1})) \\ 0, \text{ otherwise} \end{cases}$$

We have, using the language of 3.1,

$$\begin{aligned}
H^j(\mathbb{S}_{\mu'}G \otimes \mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*) &= H^j(\mathbb{S}_\mu\mathcal{R} \otimes \mathbb{S}_\alpha\mathcal{Q}^*) \otimes \mathbb{S}_{\mu'}G \\
&= H^j(\mathcal{V}(0/\mu, \alpha)) \otimes \mathbb{S}_{\mu'}G.
\end{aligned}$$

Since $\alpha_1 \leq n - r$, one easily sees that $l(\mu) = \alpha_1$. In fact, if $l(\mu) < \alpha_1$, we would get $(0, \cdots, 0, 1, \cdots)$ performing the symmetric group action described in 3.1; and we would get negative entries followed by 0s performing the symmetric group action if $l(\mu) > \alpha_1$.

Note also that we must have $\mu \supset \alpha$. Otherwise we would get 0s followed by positive entries performing the symmetric group action.

By performing adjacent transposition actions α_1 times, we can delete the first row of α, delete the first column of μ, and move everything remains in μ down by one row. Inductively, after performing $|\alpha|$ transposition actions times, we will get $(0, \cdots, 0, \alpha'/\mu', 0^{r-t})$.

To get non-trivial $H^{|\mu|-1}$, we have to perform $|\mu| - |\alpha| - 1$ transposition actions to $(\alpha'/\mu', 0^{r-t})$ to make it dominant. The only possibility is $\alpha'/\mu' = (0, \cdots, 0, -r + t + 1)$, and the corresponding $\mu = (\alpha' + (r - t, 0^{n-r-1}))$ and $H^{|\mu|-1} = \mathbb{S}_{(\alpha, 1^{r+1-t}, 0^{m-r-1})} E \otimes \mathbb{S}_{(1^{r+1-t})} F^*$.

By the same argument, one proves (2). □

8. Other examples

Here we look at some non-commutative desingularizations beyond representations with finitely many orbits. The mechanics we developed in Section 2 work in these examples. But as one have seen in Proposition 5.10, computing quiver with relations involves the problem of inner plethysm and cannot be done in general.

8.1. **Rank varieties of anti-symmetric tensors.** Let E be a vector space over k of dimension n and $H_a^{(d)}$ be the affine space $\wedge^d E^*$ consisting of anti-symmetric tensors of power d. Upon choosing a set of basis for E, it's coordinate ring is identified with $A_a^{(d)}$ $\mathrm{Sym}(\wedge^d E)$. Let $X_a^{(d)} \subset H_a^{(d)}$ be the rank variety consisting of tensors of rank $\leq n - 1$. We will find a non-commutative desingularization for $X_a^{(d)}$.

Let us review a commutative desingularization. Let

$$0 \to \mathcal{R} \to E \times \mathrm{Grass} \to \mathcal{Q} \to 0$$

be the tautological sequence over Grass where $\mathrm{Grass} = \mathrm{Grass}(1, E)$ is the grassmannian of lines in E. Let $\eta_a^{(d)}$ be the vector bundle $\wedge^d \mathcal{Q}$. As can be found in Section 7.3 of [W03], the total space $\mathcal{Z}_a^{(d)}$ of the vector bundle $\eta_a^{(d)*}$ is a commutative desingularization, i.e., $\mathcal{Z}_a^{(d)} = \underline{\mathrm{Spec}}_{\mathrm{Grass}}(\mathrm{Sym} \wedge^d \mathcal{Q})$. Alternatively, it can also be defined as the incidence variety $\mathcal{Z}_a^{(d)} = \{(S, \phi) \in \mathrm{Grass} \times H_a^{(d)} \mid \phi \in \wedge^d S \subset \wedge^d E^*\}$.

As before, let $\mathcal{Y}_a^{(d)}$ be the product $\mathrm{Grass} \times H_a^{(d)}$, p and q be the projections to Grass and $H_a^{(d)}$ respectively. Inside of $\mathcal{Y}_a^{(d)}$, there is an incidence variety, denoted by $\mathcal{Z}_a^{(d)}$. The inclusion $\mathcal{Z}_a^{(d)} \hookrightarrow \mathcal{Y}_a^{(d)}$ is denoted by j, and $X_a^{(d)} \to H_a^{(d)}$ by i. The induced map $\mathcal{Z}_a^{(d)} \to X_a^{(d)}$ from $q : \mathcal{Y}_a^{(d)} \to H_a^{(d)}$ is denoted by q' and $\mathcal{Z}_a^{(d)} \to \mathrm{Grass}$ from $p : \mathcal{Y}_a^{(d)} \to \mathrm{Grass}$ by p'.

These notations are summarized by the following diagram.

$$\begin{tikzcd}
\mathcal{Z}_a^{(d)} \arrow[dr, hook, "j"] \arrow[drr, bend left, "p'"] \arrow[dd, "q'"'] & & \\
& \mathcal{Y}_a^{(d)} = \mathrm{Grass} \times H_a^{(d)} \arrow[r, "p"] \arrow[d, "q"'] & \mathrm{Grass} = \mathrm{Grass}_1(E) \\
X_a^{(d)} \arrow[r, hook, "i"] & H_a^{(d)} = \wedge^d (E^*) &
\end{tikzcd}$$

We consider the inverse image of $\mathscr{T}il_K = \oplus_{i=0}^{n-1} \wedge^i \mathcal{Q}^*$, the Kapranov's tilting bundle over $\mathrm{Grass}(1, E)$, by $p' : \mathcal{Z}_a^{(d)} \to \mathrm{Grass}$. We can show

Lemma 8.1. *For all* $k > 0$, *and* $i, j = 0, \ldots, n-1$, $H^k(\mathcal{Z}_a^{(d)} p'^* \mathscr{H}om_{\mathcal{O}_{\mathrm{Grass}}}(\wedge^i \mathcal{Q}^*, \wedge^j \mathcal{Q}^*)) = 0$.

Proof. Use the Cauchy-Littlewood and the Littlewod-Richardson formulas, we can reduce the sheaf $p'^{*}\mathscr{H}om_{\mathscr{O}_{\text{Grass}}}(\wedge^i Q^*, \wedge^j Q^*) \cong \mathscr{H}om_{\mathscr{O}_{\text{Grass}}}(\wedge^i Q^*, \wedge^j Q^* \otimes_{\mathscr{O}_{\text{Grass}}} \text{Sym}(\wedge^d Q))$ into the form

$$\oplus(\mathbb{S}_\gamma Q^*)^{\oplus C_\gamma}$$

for some coefficients C_γ. By the Bott's theorem 3.2, all the higher cohomology of

$$\mathscr{H}om_{\mathscr{O}_{\text{Grass}}}(\wedge^i Q^*, \wedge^j Q^* \otimes_{\mathscr{O}_{\text{Grass}}} \text{Sym}(\wedge^d Q))$$

vanishes. □

From the Lemma above, using Theorem 2.7, we get the following

Proposition 8.2. *The Kapranov's tilting bundle over* $\text{Grass}(1, E)$ *is denoted by* $\mathscr{T}il_K$. *The rank* $n-1$ *subvariety* $X_a^{(d)}$ *of d-th anti-symmetric tensors and its desingularization* $\mathcal{Z}_a^{(d)}$ *are as above. The bundle* $p'^{*}\mathscr{T}il_K$ *is a tilting bundle over* $\mathcal{Z}_a^{(d)}$.

Since $\mathcal{Z}_a^{(d)}$ is smooth, $\text{End}_{\mathcal{Z}_a^{(d)}}(p'^{*}\mathscr{T}il_K)$ has finite global dimension.

Lemma 8.3. *Notations as above,* $\text{End}_{\mathcal{Z}_a^{(d)}}(p'^{*}\mathscr{T}il_K)$ *is maximal Cohen-Macaulay iff* $\binom{n-2}{d-1} - n - 1 \geq 0$.

Remark 8.4. As can be easily checked, $\binom{n-2}{d-1} - n - 1 \geq 0$ if and only if $n \geq 6$ and $3 \leq d \leq n-3$.

Proof. By Theorem 3.4, it suffices to compute $H^k(\text{Grass}, \wedge^i Q \otimes \wedge^j Q^* \otimes \omega_{\text{Grass}} \otimes \wedge^{top}\xi^* \otimes \text{Sym}(\wedge^d Q))$ for $k > 0$ and $i, j = 0, \ldots, n-1$, here $\xi = \mathcal{R} \otimes \wedge^{d-1} Q$ according to 7.3.1 of [W03]. The sheaf $\omega_{\text{Grass}} = \wedge^{n-1} Q^* \otimes \mathcal{R}^{n-1}$. Using Cauchy-Littlewood and Littlewood-Richardson formulas, $\wedge^i Q \otimes \wedge^j Q^* \otimes \omega_{\text{Grass}} \otimes \wedge^{top}\xi^* \otimes \text{Sym}(\wedge^d Q) \cong \wedge^{j-i} Q^* \otimes \text{Sym}\wedge^d Q \otimes \wedge^{n-1} Q^{\binom{n-2}{d-1}-(n-1)-1} \otimes \wedge^n E^{*\binom{n-1}{d-1}-(n-1)}$. The conclusion now follows from the Bott's Theorem. □

We know that $\mathscr{O}_{\text{Grass}}$ is a direct summand of $\mathscr{T}il_K$. In the case $\binom{n-2}{d-1} - n - 1 \geq 0$, the module $q'_* p'^{*}\mathscr{T}il_K$ is maximal Cohen-Macaulay. In particular, it is reflexive. According to Proposition 2.7 and Lemma 2.8, using Lemma 8.9, we get the following Proposition.

Proposition 8.5. *Notations as above, in the case* $\binom{n-2}{d-1} - n - 1 \geq 0$, *the map*

$$\text{End}_{\mathcal{Z}_a^{(d)}}(p'^{*}\mathscr{T}il_K) \to \text{End}_{A_a^{(d)}}(q'_* p'^{*}\mathscr{T}il_K)$$

is an isomorphism of rings and $\text{End}_{\mathcal{Z}_a^{(d)}}(p'^{*}\mathscr{T}il_K)$ *is a non-commutative crepant desingularization of* $X_a^{(d)}$.

Remark 8.6. In this case, the tilting bundle over $\mathcal{Z}_a^{(d)}$ is the inverse image of an exceptional collection $\{\wedge^i Q^* \mid i = 0, \ldots, n-1\}$ over Grass. As can be checked by definition, The dual collection is given by $\wedge^{n-1} Q^* \otimes \mathbb{S}_{n-i}\mathcal{R}[n-1-i]$.

Let us take n=6 and d=3, for an example. One can prove as in Proposition 5.10, that for any $\alpha, \beta = 0, \ldots, 6$, $\text{Ext}^t(S_\alpha, S_\beta) = H^{t-s-\beta+\alpha}(\text{Grass}, \wedge^s \wedge^3 Q^* \otimes \mathbb{S}_{(6-\beta)}\mathcal{R} \otimes \mathbb{S}_{(6-\alpha)}\mathcal{R}^*)$. As it is known that $\wedge^2 \wedge^3 \mathbb{C}^6 \cong \mathbb{S}_{2,2,1,1,0,0}\mathbb{C}^6 \oplus \mathbb{S}_{1,1,1,1,1,1}\mathbb{C}^6$. This means, in this example Ext^1 and Ext^2 can be calculated very easily which gives the quiver with relations for the endomorphism algebra.



and the relations:

$$\alpha_i\alpha_{i+1}(\wedge^2 E);$$
$$\beta\alpha\beta\alpha + \alpha\beta\alpha\beta(\wedge^2 E).$$

8.2. **Cone over a rational normal curve.** Let E be a vector spaces over k of dimension n and $H_s^{(d)}$ be the affine space $\mathbb{S}_d E^*$ consisting of symmetric tensors of power d. Upon choosing a set of basis for E, it's coordinate ring is identified with $A_s^{(d)} = \mathrm{Sym}(\mathbb{S}_d E)$. Let $X_s^{(d)} \subset H_s^{(d)}$ be the rank variety consisting of symmetric tensors of rank $\leq n-1$. We will find a non-commutative desingularization for $X_s^{(d)}$ and describe it in certain cases.

Let us review a commutative desingularization. Let

$$0 \to \mathcal{R} \to E \times \mathrm{Grass} \to \mathcal{Q} \to 0$$

be the tautological sequence over Grass where Grass $= \mathrm{Grass}(1, E)$ is the grassmannian of lines in E. Let $\eta_s^{(d)}$ be the vector bundle $\mathbb{S}_d\mathcal{Q}$. As can be found in Section 7.3 of [W03], the total space $\mathcal{Z}_s^{(d)}$ of the vector bundle $(\eta_s^{(d)})^*$ is a commutative desingularization, i.e., $\mathcal{Z}_s^{(d)} = \underline{\mathrm{Spec}}_{\mathrm{Grass}}(\mathrm{Sym}\,\mathbb{S}_d\mathcal{Q})$. Alternatively, it can also be defined as the incidence variety $\mathcal{Z}_s^{(d)} = \{(S, \phi) \in \mathrm{Grass} \times H_s^{(d)} \mid \phi \in \mathbb{S}_d S \subset \mathbb{S}_d E^*\}$.

As before, let $\mathcal{Y}_s^{(d)}$ be the product Grass $\times H_s^{(d)}$, p and q be the projections to Grass and $H_s^{(d)}$ respectively. Inside of $\mathcal{Y}_s^{(d)}$, there is an incidence variety, denoted by $\mathcal{Z}_s^{(d)}$. The inclusion $\mathcal{Z}_s^{(d)} \hookrightarrow \mathcal{Y}_s^{(d)}$ is denoted by j, and $X_s^{(d)} \to H_s^{(d)}$ by i. The induced map $\mathcal{Z}_s^{(d)} \to X_s^{(d)}$ from $q : \mathcal{Y}_s^{(d)} \to H_s^{(d)}$ is denoted by q' and $\mathcal{Z}_s^{(d)} \to$ Grass from $p : \mathcal{Y}_s^{(d)} \to$ Grass by p'.

These notations are summarized by the following diagram.

$$\begin{tikzcd}
\mathcal{Z}_s^{(d)} \arrow[dr, hook, "j"] \arrow[drr, bend left, "p'"] \arrow[dd, "q'"'] & & \\
 & \mathcal{Y}_s^{(d)} = \mathrm{Grass} \times H_s^{(d)} \arrow[r, "p"] \arrow[d, "q"'] & \mathrm{Grass} = \mathrm{Grass}_1(E) \\
X_s^{(d)} \arrow[r, hook, "i"] & H_s^{(d)} = \mathbb{S}_d(E^*) &
\end{tikzcd}$$

As before, we have:

Lemma 8.7. *For all* $k > 0$, *and* $i, j = 0, \ldots, n-1$, $H^k(\mathcal{Z}_s^{(d)} p'^* \mathscr{H}\mathrm{om}_{\mathscr{O}_{\mathrm{Grass}}}(\wedge^i \mathcal{Q}^*, \wedge^j \mathcal{Q}^*)) = 0$.

From the Lemma above, using Theorem 2.7, we get the following

Proposition 8.8. *The Kapranov's tilting bundle over* Grass$(1, E)$ *is denoted by* $\mathscr{T}il_K$. *The rank* $n-1$ *subvariety* $X_s^{(d)}$ *of d-th symmetric tensors and its desingularization* $\mathcal{Z}_s^{(d)}$ *are as above. The bundle* $p'^*\mathscr{T}il_K$ *is a tilting bundle over* $\mathcal{Z}_s^{(d)}$.

Since $\mathcal{Z}_s^{(d)}$ is smooth, $\mathrm{End}_{\mathcal{Z}_s^{(d)}}(p'^*\mathscr{T}il_K)$ has finite global dimension.

Lemma 8.9. *Notations as above,* $\mathrm{End}_{\mathcal{Z}_s^{(d)}}(p'^*\mathscr{T}il_K)$ *is maximal Cohen-Macaulay.*

Proof. By Theorem 3.4, it suffices to compute $H^k(\mathrm{Grass}, \wedge^i\mathcal{Q} \otimes \wedge^j\mathcal{Q}^* \otimes \omega_{\mathrm{Grass}} \otimes \wedge^{top}\xi^* \otimes \mathrm{Sym}(\mathbb{S}_d\mathcal{Q}))$ for $k > 0$ and $i, j = 0, \ldots, n-1$, here $\xi = \mathcal{R} \otimes \mathbb{S}_{d-1}E$ according to 7.2.1 of [W03]. The sheaf $\omega_{\mathrm{Grass}} = \wedge^{n-1}\mathcal{Q}^* \otimes \mathcal{R}^{n-1}$. Using Cauchy-Littlewood and Littlewood-Richardson formulas, $\wedge^i\mathcal{Q} \otimes \wedge^j\mathcal{Q}^* \otimes \omega_{\mathrm{Grass}} \otimes \wedge^{top}\xi^* \otimes \mathrm{Sym}(\mathbb{S}_d\mathcal{Q}) \cong \wedge^{j-i}\mathcal{Q}^* \otimes \mathrm{Sym}\,\mathbb{S}_d\mathcal{Q} \otimes \wedge^{n-1}\mathcal{Q}^{\binom{n+d-1}{n-1}-(n-1)-1} \otimes \mathbb{S}_\mu E$ for some μ. Now it follows easily from the Bott's Theorem that $\mathrm{End}_{\mathcal{Z}_s^{(d)}}(p'^*\mathscr{T}il_K)$ is maximal Cohen-Macaulay iff $\binom{n+d-1}{n-1} - n - 1 \geq 0$. But as can be easily seen, $\binom{n+d-1}{n-1} - n - 1$ is always positive. $\square$

In the case that $n = 2$, one can see that $X_s^{(d)}$ is the cone over a rational normal curve. And in this case, $\mathrm{End}_{\mathcal{Z}_s^{(d)}}(p'^* \mathscr{T}il_K)$ is always maximal Cohen-Macaulay grant that $d \geq 2$.

With similar calculation as in Proposition 5.10, we get the following:

Proposition 8.10. *For any two simple objects S_i, S_j, $i, j = 0, 1$, we have*

$$\mathrm{Ext}^t(S_i, S_j) = \oplus_{s=0}^{1} H^{t-s-j+i}(\mathbb{P}^1, (Q^*)^{ds} \otimes (\mathcal{R}^*)^{j-i}).$$

Plug-in $t = 1, 2$ and $i, j = 0, 1$, we get the quiver with relations for the non-commutative desingularization.

$$\begin{tikzcd} \bullet_0 \arrow[r, bend left, "\beta(\mathbb{S}_{d-1}E)"] & \bullet_1 \arrow[l, bend left, "\alpha(E)"] \end{tikzcd}$$

and the relations:

$$\alpha\beta(\mathbb{S}_{(d-1,1)}E);$$
$$\beta\alpha(\mathbb{S}_{(d-1,1)}E).$$

Remark 8.11. Many other examples of representations with finitely many orbits will be analyzed in subsequent papers.

References

[ASS06] I. Assem, D. Simson, and A. Skowronski, *Elements of the Representation Theory of Associative Algebras Volume 1 Techniques of Representation Theory*, Lond. Math. Soc. Student Texts, vol. 65 Cambridge University Press, New York, 2006. MR2197389 2.4, 2.4

[Be06] R. Bezrukavnikov, *Noncommutative counterparts of the Springer resolution*, Proceedings of the International Congress of Mathematicians, vol. II, Eur. Math. Soc., Zürich, 2006, pp. 1119-1144. MR2275638 2.3, 2.28

[Br05] T. Bridgeland, *t-structures on some local Calabi-Yau varieties*, preprint, `arXiv:0502050v1`, 2005. 1

[BH10] R. Buchweitz, L. Hille, *Hoschild cohohomology of schemes with tilting object*, preprint, `arXiv:1003.4201`, 2010. 1

[Bou98] N. Bourbaki, *Commutative algebra*, Chapters 1-7, Reprint of the 1989 English translation, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1998.MR1727221 2.2, 2.2

[BO02] A. Bondal, D. Orlov, *Derived Categories of Coherent Sheaves*, Proceedings of the International Congress of Mathematicians, Vol. II, Beijing, 2002, pp. 47-56. `arXiv:0206295`, 2002.MR1957019 2.2, 2.3

[BLV09] R. Buchweitz, G. Leuschke, M. van den Bergh, *non-commutative desingularization of determinantal varieties 1*, Invent. Math. 182 (2010), no. 1, pp. 47-115. `arXiv:0911.2659v2`MR2672281 1, 1, 1, 3.2, 4.7, 7.1

[BLV11] R. Buchweitz, G. Leuschke, M. van den Bergh, *non-commutative desingularization of determinantal varieties 2*, preprint, `arXiv:1106.1833v1`, 2011. 1, 2.2, 7.2

[CG00] N. Chriss, V. Ginzburg, *Representation Theory and Complex Geometry*, 2nd ed., Birkhäuser, Boston, 2000. MR1433132 2.4

[F97] W. Fulton, *Young Tableaux*, Lond. Math Soc. Student Texts, vol. 35, Cambridge University Press, New York, 1997. MR1464693 1, 3.1

[GM03] S. Gelfand, Y. Manin, *Methods of Homological Algebra* 2nd ed, Springer-Verlag, Berlin-Heidelberg, 2003. MR1950475 2.2, 2.28

[H77] R. Hartshorne, *Algebraic geometry*, Grad. Texts in Math., vol. 52. Springer-Verlag, New York-Heidelberg, 1977. MR0463157 2.4

[HV07] L. Hille, M. van den Bergh, *Fourier-Mukai Transforms, Handbook of Tilting Theory*, London Math. Soc. Lecture Note Ser. vol. 332, Cambridge University Press, 2007, 147-77. MR2384610 1, 2.1, 2.10, 5.2, 6.2, 7.2

[K88] M.M. Kapranov, *on the derived categories of coherent sheaves on some homogeneous spaces*, Invent. Math. **92** (1988), no.3, 479-509. MR939472 3.2

[SW10] S.V. Sam, J. Weyman, *Pieri resolutions for classical groups*, preprints, `arXiv:0907.4505v4` 2010. 5.5

[V04a] M. van den Bergh, *three dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no.3, pp. 423-445. MR2057015 2.2

[V04b] M. van den Bergh, *non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749-770. `arXiv:0211064`MR2077594 1

[W03] J. Weyman, *Cohomology of Vector Bundles and Syzygies*, Cambridge Tracts in Math., vol. 149. Cambridge University Press, New York, 2003. MR1988690 1, 1, 1, 2.2, 3.1, 3.3, 3.1, 5.1, 5.2, 6.1, 6.2, 7.1, 8.1, 8.1, 8.2, 8.2

Department of Mathematics, Northeastern University, Boston, MA 02115, USA
E-mail address: `j.weyman@neu.edu`

Department of Mathematics, Northeastern University, Boston, MA 02115, USA
E-mail address: `zhao.g@husky.neu.edu`